

Crown Copyright 2002 First Published 2002

Printed in the United Kingdom

Enquiries regarding this report should be directed to:

Head of Woodland Surveys Forest Research Forestry Commission 231 Corstorphine Road Edinburgh EH12 7AT

Telephone: 0131 314 6122

Email: woodland.surveys@forestry.gsi.gov.uk

CONTENTS

| Acknowle | dgements | V |
|---|--|--|
| Introductio | on | 1 |
| Backgrour Survey me Main point Inventory I | thod ts from the survey results | 1 1 2 2 |
| Map 1: Map 2: Map 3: Map 4: | County boundaries Distribution of woodland over 2 hectares Distribution of woodland over 2 hectares by ownership Distribution of woodland over 2 hectares by Interpreted Forest Type | 3 4 5 6 |
| Summary i | results from the National Inventory of Woodland and Trees (NIWT) | 7 |
| Tables 1 – | 5 | |
| Table 1: Table 2: Table 3: Table 4: Table 5: | Woodland area by woodland size class Woodland area by forest type and woodland size Woodland area by principal species and woodland size Numbers of live trees outside woodland by feature type Lengths of Linear Features | 9 10 11 12 12 |
| Results from | m the Main Woodland Survey (MWS) | 13 |
| Tables 6 - 1 | 12 | |
| Table 6: Chart: Table 7a: Table 8: Chart: Table 9a: Graph: Table 9b: Graph: | Summary of woodland area by ownership Woodland area by ownership Size class distribution of woodland Size class distribution of woodland by ownership units Area of woodland by forest type and ownership Area of woodland by forest type Area of High Forest by principal species and ownership Area of High Forest by principal species and ownership Area of High Forest by principal species, ownership and category High Forest Category 1 Area by principal species and ownership High Forest Category 2 | 15 15 16 16 17 17 18 19 20 |
| Table 10a: | Area by principal species and ownership High Forest Category 1 | 21 |
| Graph: | Area by principal species and planting year class High Forest Category 1 Area by planting year class | 22 23 |

NATIONAL INVENTORY OF WOODLAND AND TREES - CORNWALL

| Table 10b: | High Forest Category 1 | 0.4 |
|------------------------|---|----------|
| Graph: | Forestry Commission: area by principal species and planting year class High Forest Category 1 | 24 |
| | Forestry Commission - area by planting year class | 25 |
| Table 10c: | High Forest Category 1 Other ownership : area by principal species and planting year class | 26 |
| Graph: | High Forest Category 1 | 20 |
| · | Other ownership: area by planting year class | 27 |
| Table 11: Table 12: | High Forest: principal species by planting year class Ownership type by area and percentage | 28 29 |
| Chart: | Ownership type by area | 29 |
| Dogwille from | the Company of Creat Mean district and Trace (CCMT) | 31 |
| Results Itom | the Survey of Small Woodland and Trees (SSWT) | 31 |
| Tables 13 – 1 | 18 | |
| Table 13: | Summary of information from the Survey of Small Woodland and Trees | 33 |
| Table 14: | Woodland area by feature type and woodland size | 33 |
| Table 15: Table 16: | Numbers of live trees outside woodland by species and feature type Numbers of dead trees outside woodland by species and feature type | 34 35 |
| Table 18: | Numbers of dead frees outside woodland by species and height band | 36 |
| Table 18: | Numbers of Groups by group size | 37 |
| Comparisor | of results with the 1980 Census and previous surveys | 39 |
| Tables 19 - : | 23 | |
| Table 19: | Comparison of woodland area between 1980 Census and 1997 Inventory | 41 |
| Table 20: | Comparison of High Forest area by species between 1980 Census and 1997 Inventory | 42 |
| Chart: | Comparison of High Forest Carlo your Large by species between 1980 Census and 1997 Inventory | 43 |
| Table 21: | Comparison of High Forest Category 1 area by planting year class between 1980 Census and 1997 Inventory | 44 |
| Chart: | Comparison of High Forest Category 1 area by planting year class | 45 |
| | between 1980 Census and 1997 Inventory | |
| Table 22: Table 23: | Comparison of numbers of live trees outside woodland between 1980 Census and 1997 Inventory Comparison of density of non-woodland features between 1980 Census and 1997 Inventory | 46 46 |
| | | 40 |
| Woodland o | cover Change in woodland cover through time (1890 – 2000) | 47 |
| Maps: | Woodland cover hirough time (1895 – 1998) | 47 |
| Glossary | | 49 |
| Glossury | | 47 |

ACKNOWLEDGEMENTS

The Forestry Commission is grateful to many people who helped in the completion of this survey. In particular, the Forestry Commission would like to thank owners and occupiers of the land selected for sampling.

Woodland Surveys Branch of Forest Research was responsible for carrying out the survey and analysing the data. A large number of Forestry Commission and contract staff were involved in the survey from its inception.

Preparation of the digital cartography for Cornwall was carried out by Graham Bull, Woodland Survey Officer, and Woodland GIS Officers Chris Brown, Robert Beck and Esther Whitton. Data processing and analysis was carried out by Woodland Data Officers Justin Gilbert and Shona Cameron.

The authors of this Report are Steve Smith (Head of Woodland Surveys) and Justin Gilbert (Woodland Data Officer) of Forest Research.

| NATIONAL INVENTORY OF WOODLAND AND TREES – CORNWALL | | |
|---|----|--|
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | vi | |

INTRODUCTION

This report presents the results for Cornwall from the Forestry Commission National Inventory of Woodland and Trees (NIWT).

The Inventory consists of two separate surveys -

- The Main Woodland Survey (MWS) covering woodlands of 2 hectares and over
- The Survey of Small Woodland and Trees (SSWT) covering Small Woods, Groups of Trees, Linear Features and Individual Trees.

BACKGROUND

Since 1924 the Forestry Commission has carried out a number of national woodland surveys at intervals of between 15 and 20 years. The previous survey was carried out between 1979 and 1982. With the statistics becoming increasingly out of date the Forestry Commission decided to undertake a new survey: the *National Inventory of Woodland and Trees*.

The survey fieldwork for Great Britain was completed in July 2000. Work began in Scotland in 1994, followed by Southern England, Wales and Northern England.

SURVEY METHODS

Main Woodland Survey

In England, Woodland Surveys derived a digital map of all woodland showing Interpreted Forest Types from 1:25 000 scale aerial photography. This provided the basis for the sampling.

The digital map gives the extent of all woodland over 2 hectares and this was updated as survey work progressed. The maps on pages 4-6 show: overall woodland cover; woodland by ownership; and woodland by Interpreted Forest Type, respectively. The total area of woodland was obtained from the digital map with ground sampling undertaken to evaluate a wide range of woodland information such as species, age and stocking.

From the digital map the area of each woodland was recorded and this information was used to determine the intensity at which any selected woodland would be sampled. The overall sampling scheme was as follows:

2.0ha - <100ha : every fifth wood
 100ha - <500ha : two woods in five

500ha and larger : all woods

1 hectare square plots were used to sample the selected woodlands on the ground. This was a change of practice from all previous Census surveys, where whole woods have been selected for survey. For each of the three bands of woodland area a different sampling grid was used with the density of the squares being reduced as the woodlands increase in size. The overall aim was to sample 1% of the woodland in each size class.

Survey of Small Woodland and Trees

The land area of England was stratified into coastal and inland 1 km x 1 km squares and a random sample of 1 km² plots were then selected, representing around 1% of the land area. 1:25 000 scale aerial photos were then used to identify features in each sample square. Each 1 km² was then divided into 16 parts, and two of these were selected at random for field data collection. Data was collected on Small Woodlands (0.10 - <2.00 ha), Linear Features, Groups and Individual Trees. The survey did not collect information from areas of developed land of 2 hectares or more.

MAIN POINTS FROM THE SURVEY RESULTS


- The total area of woodland of 0.1 hectares and over in Cornwall is 26,869 hectares. This represents 7.5 % of the land area. (Table 1)
- Broadleaved woodland is the dominant forest type representing 66.1 % of all woodland. Conifer woodland represents 19.0 %, Mixed woodland 10.2 % and Open Space within woodlands 2.4 %. (Table 2)
- The main conifer species is Sitka spruce covering 1,140 hectares or 18.8 % of all conifer species. The main broadleaved species is Oak covering 2,395 hectares or 12.3 % of all broadleaved species. (Table 3)
- 3,035 hectares or 11 % of woodland over 2 hectares is owned by or leased to the Forestry Commission, and 23,736 hectares or 89 % of woodland is in Other ownership. (Table 6)
- There are a total of 1,675 woods over 2 ha within Cornwall with a mean wood area of 16.1 hectares. (Table 7a) There are a total of 164 woods from 0.1 <2.0 hectares with a mean wood area of 0.60 hectares. (Table 14)
- There are 1.3 million live trees outside woodland in Cornwall. (Table 15)
- Woodland land cover increased by over 8,500 hectares from 5.1 % to 7.5 % of the land area between 1980 and 1997. (Table 19)
- The area of broadleaves increased by 68% between 1980 and 1997, with the relative proportion of broadleaves to conifers increasing from 69% to 76%. (Table 20)

INVENTORY REPORTS


As well as this report for Cornwall, reports are available for the other counties in the region as shown on the map opposite. Also available are region and county reports for England as well as a report for the country as a whole. Wales and Scotland are also covered by reports.




Map 2 Distribution of woodland over 2 hectares



Map 3 Distribution of woodland over 2 hectares by ownership



Map 4 Distribution of woodland over 2 hectares by Interpreted Forest Type



SUMMARY RESULTS FROM THE NATIONAL INVENTORY OF WOODLAND AND TREES (NIWT)

Both the Main Woodland Survey and the Survey of Small Woodland and Trees contributed to the estimate of woodland area for Cornwall.

Tables 1-3 show the combined woodland area from the Main Woodland Survey and the Survey of Small Woodland and Trees.

Tables 4 and 5 summarise the numbers of live trees outside woodland, and the lengths of Linear Features from the Survey of Small Woodland and Trees.

Table 1: Woodland area by woodland size class

Table 2: Woodland area by forest type and woodland size
 Table 3: Woodland area by principal species and woodland size
 Table 4: Numbers of live trees outside woodland by feature type

Table 5: Lengths of Linear Features

Note: The figures in many of the tables may not add due to rounding



Table 1 Woodland area by woodland size class

| Woodland size (ha) | Woodland area (ha) | % of Woodland area |
|------------------------|--------------------|--------------------|
| 2.00 and over | 26,771 | 99.6 |
| 0.25 - < 2.00 | 93 | 0.3 |
| 0.10 - < 0.25 | 5 | 0.0 |
| Total area of woodland | 26,869 | 100.0 |
| % Woodland land cover | 7.5 | |

^{1.} Area of Cornwall, including inland water, 356,466 ha based on digital boundaries used in the 1991 Census of Population

Table 2 Woodland area by forest type and woodland size

| Forest type | Woodland 2.0 and over | l size (ha) 0.1 - <2.0 | Total area (ha) | Percentage of total area |
|------------------|--------------------------|---------------------------|--------------------|--------------------------|
| Conifer | 5,095 | 1 | 5,096 | 19.0 |
| Broadleaved | 17,694 | 64 | 17,758 | 66.1 |
| Mixed | 2,706 | 33 | 2,739 | 10.2 |
| Coppiced | 291 | 0 | 291 | 1.1 |
| Copp-w-standards | 271 | 0 | 271 | 1.0 |
| Windblow | 0 | 0 | 0 | 0.0 |
| Felled | 59 | 0 | 59 | 0.2 |
| Open Space | 656 | 1 | 657 | 2.4 |
| Total | 26,771 | 98 | 26,869 | 100 |

See Glossary for definitions of forest types.

Table 3 Woodland area by principal species and woodland size

| Species/Groups | Woodland size (ha) | | Total area | Percentage | of total area |
|----------------------|--------------------|-----------|------------|------------|---------------|
| | 2.0 and over | 0.1 -<2.0 | (ha) | Category* | Species** |
| Pine | 199 | 1 | 200 | 3.3 | 0.8 |
| Sitka spruce | 1,140 | 0 | 1,140 | 18.8 | 4.5 |
| Larch | 734 | 8 | 742 | 12.2 | 2.9 |
| Other conifers | 2,925 | 4 | 2,929 | 48.2 | 11.4 |
| Mixed conifers | 1,064 | 0 | 1,064 | 17.5 | 4.2 |
| Total conifers | 6,062 | 13 | 6,075 | 100.0 | 23.7 |
| Oak | 2,386 | 9 | 2,395 | 12.3 | 9.4 |
| Beech | 576 | 30 | 606 | 3.1 | 2.4 |
| Sycamore | 569 | 14 | 583 | 3.0 | 2.3 |
| Ash | 652 | 10 | 662 | 3.4 | 2.6 |
| Birch | 211 | 0 | 211 | 1.1 | 0.8 |
| Elm | 0 | 4 | 4 | 0.0 | 0.0 |
| Other broadleaves | 1,485 | 16 | 1,501 | 7.7 | 5.9 |
| Mixed broadleaves | 13,556 | 2 | 13,558 | 69.5 | 53.0 |
| Total broadleaves | 19,433 | 84 | 19,518 | 100.0 | 76.3 |
| Total all species*** | 25,496 | 97 | 25,593 | | 100.0 |

^{*}Category - species/group percentage of conifer or broadleaved category **Species/group percentage of all species

The standard errors of the area estimates for woodland of 2 ha and over tor the most common species or species groups are as tollows

| Coniters | /% |
|--------------|-----|
| Broadleaves | 3% |
| Sitka spruce | 22% |
| Oak | 14% |
| Ash | 18% |
| | |

Where the standard errors of these summary measures are 10% or less, the confidence intervals will be approximately symmetrical; the true value is expected to be within +/- one standard error for about 68% (or about two-thirds) of all cases, and within +/- two standard errors for about 95% of all cases. Where percentage standard errors are larger, e.g. for less common species or more variable species composition, the confidence intervals will be less symmetrical (and wider).

11

^{***}Excludes the 1,278 ha of Coppice, Felled and Open space areas which were included in Table 2

Table 4 Numbers of live trees outside woodland by feature type

| Feature type | Total number of features | Total number of live trees | Mean number of trees per feature | Tree density (per sq km) |
|------------------------|-----------------------------|-------------------------------|-------------------------------------|-----------------------------|
| Groups | 96,400 | 1,022,400 | 11 | 287 |
| Narrow Linear Features | 4,900 | 186,700 | 38 | 52 |
| Individual Trees | 59,600 | 59,600 | 1 | 17 |
| Total | | 1,268,700 | | 356 |

- 1. Land area used to calculate tree density 356,466 ha based on digital boundaries used in 1991 Census of Population
- 2. The standard errors of the live tree number estimates for these feature types are:

| Groups | 26% |
|------------------------|-----|
| Narrow Linear Features | 58% |
| Individual Trees | 24% |

- 3. Where the standard errors of these summary measures are 10% or less, the confidence intervals will be approximately symmetrical; the true value is expected to be within +/- one standard error for about 68% (or about two-thirds) of all cases, and within +/- two standard errors for about 95% of all cases. Where percentage standard errors are larger, e.g. for less common species or more variable species composition, the confidence intervals will be less symmetrical (and wider).
- 4. See Glossary for definitions of feature types.

Table 5 Lengths of Linear Features

| Feature type | Total number of features | Total length of features (km) | Density of features (m per sq km) |
|------------------------|-----------------------------|-------------------------------|--------------------------------------|
| Wide Linear Features | 0 | 0 | 0 |
| Narrow Linear Features | 4,900 | 295 | 83 |
| Total | | 295 | 83 |

- 1. Land area used to calculate tree density 356,466 ha based on digital boundaries used in 1991 Census of Population
- 2. The standard errors of the length estimates for these feature types are:

Wide Linear Features Narrow Linear Features - 67%

- 3. Where the standard errors of these summary measures are 10% or less, the confidence intervals will be approximately symmetrical; the true value is expected to be within +/- one standard error for about 68% (or about two-thirds) of all cases, and within +/- two standard errors for about 95% of all cases. Where percentage standard errors are larger, e.g. for less common species or more variable species composition, the confidence intervals will be less symmetrical (and wider).
- 4. See Glossary for definitions of feature type .

RESULTS FROM THE MAIN **WOODLAND SURVEY (MWS)**

Survey Method

Woods were selected from the digital map of woodland of 2 hectares and over, then sampled using a random grid of 1 hectare sample plots. The density of sample plots was reduced as the sampled woodland increase in size, the general aim being to sample 1% of the woodland area. The ground sampling evaluated a wide range of data such as species, age and stocking.

Table 6: Summary of woodland area by ownership

Chart: Woodland area by ownership Table 7a: Size class distribution of woodland

Table 7b: Size class distribution of woodland by ownership units Table 8: Area of woodland by forest type and ownership

Chart: Area of woodland by forest type

Area of High Forest by principal species and ownership Table 9a: Area of High Forest by principal species and ownership Graph:

Table 9b: Area of High Forest by principal species, ownership and category

High Forest Category 1 Graph:

Area by principal species and ownership

Graph: High Forest Category 2

Area by principal species and ownership

Table 10a: High Forest Category 1

Area by principal species and planting year class

Graph: High Forest Category 1

Area by planting year class

Table 10b: High Forest Category 1

Forestry Commission: area by principal species and planting year class

Graph: High Forest Category 1

Forestry Commission - area by planting year class

Table 10c: High Forest Category 1

Other ownership: area by principal species and planting year class

Graph: High Forest Category 1

Other ownership: area by planting year class

Table 11: High Forest: principal species by planting year class

Table 12: Ownership type by area and percentage

Chart: Ownership type by area

Note: The figures in many of the tables may not add due to rounding



Table 6 Summary of woodland area by ownership

| Ownership | ha | % woodland |
|------------------------|--------|------------|
| Forestry Commission | 3,035 | 11 |
| Other | 23,736 | 89 |
| Total area of woodland | 26,771 | 100 |

- 1. Woodland area from aerial photographic interpretation map updated to 31 March 1997
- 2. See Glossary for definitions of ownership types




Table 7a Size class distribution of woodland

| Size class (ha) | Number of woods | Total area (ha) | Percent of total area | Mean wood area (ha) |
|--------------------|--------------------|--------------------|--------------------------|------------------------|
| <10 | 1,227 | 5,507 | 20 | 4.5 |
| 10 - <20 | 206 | 2,860 | 11 | 13.9 |
| 20 - <50 | 149 | 4,654 | 17 | 31.2 |
| 50 - <100 | 54 | 3,829 | 14 | 70.9 |
| <100 | 1,636 | 16,850 | 63 | 10.3 |
| 100 - <500 | 37 | 8,259 | 31 | 223.2 |
| 500 and > | 2 | 1,829 | 7 | 914.5 |
| All woods | 1,675 | 26,939 | 100 | 16.1 |

Table 7b Size class distribution of woodland by ownership units

| Size class (ha) | FC or Other | Number of woods | Total area (ha) | Percent of total area | Mean wood area (ha) |
|--------------------|----------------|--------------------|--------------------|--------------------------|------------------------|
| <10 | FC | 12 | 63 | 0 | 5.2 |
| | 0 | 1,322 | 5,771 | 21 | 4.4 |
| 10 - <20 | FC | 8 | 115 | 0 | 14.4 |
| | 0 | 212 | 2,962 | 11 | 14.0 |
| 20 - <50 | FC | 11 | 361 | 1 | 32.9 |
| | 0 | 159 | 4,932 | 18 | 31.0 |
| 50 - <100 | FC | 5 | 315 | 1 | 62.9 |
| | 0 | 59 | 4,162 | 15 | 70.5 |
| <100 | FC | 36 | 854 | 3 | 23.7 |
| | 0 | 1,752 | 17,827 | 66 | 10.2 |
| 100 - <500 | FC | 13 | 2,180 | 8 | 167.7 |
| | O | 27 | 6,077 | 23 | 225.1 |
| 500 and > | FC | 0 | 0 | 0 | 0.0 |
| | 0 | 0 | 0 | 0 | 0.0 |
| Total | FC | 49 | 3,035 | 11 | 61.9 |
| | 0 | 1,779 | 23,904 | 89 | 13.4 |

- Table 7a and 7b are based solely on the digital woodland map. The other MWS tables are derived from the field sample data
- 2. The total area in Tables 7a and 7b is 168 hectares more than recorded in Table 6. This is mainly due to the field samples recording some land in other land uses not differentiated from woodland in the digital map
- 3. The data available from the digital map enable the identification of woodlands according to their ownerships, Forestry Commission or Other. The entries in table 7b cannot be added to derive table 7a as some woods may consist of both Forestry Commission and Other ownership(s)

For example, the Forestry Commission may own most of a large wood with some parts in Other ownership(s). In Table 7a the whole area would be treated as one wood and the area allocated to one size category. In Table 7b each of the ownership units would be allocated to the size category for that unit. Dividing woods by ownership can occasionally generate part woods of less than 2 hectares

 Table 8
 Area of woodland by forest type and ownership

| Forest type | Forestry C | ommission | Otl | ner | All owr | nerships |
|-------------|------------|-----------|--------|-------|---------|----------|
| | ha | % | ha | % | ha | % |
| Conifer | 2,072 | 68.3 | 3,024 | 12.7 | 5,095 | 19.0 |
| Broadleaved | 493 | 16.2 | 17,201 | 72.5 | 17,694 | 66.1 |
| Mixed | 324 | 10.7 | 2,382 | 10.0 | 2,706 | 10.1 |
| Coppice | 0 | 0.0 | 291 | 1.2 | 291 | 1.1 |
| Copp-w-Stds | 0 | 0.0 | 271 | 1.1 | 271 | 1.0 |
| Windblow | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| Felled | 21 | 0.7 | 38 | 0.2 | 59 | 0.2 |
| Open Space | 126 | 4.2 | 530 | 2.2 | 656 | 2.5 |
| Total | 3,035 | 100.0 | 23,736 | 100.0 | 26,771 | 100.0 |




Table 9a Area of High Forest by principal species and ownership

| Species | Forestry | Commiss | sion | C | other | | All ow | nerships/ | |
|---------------------|----------|---------|-------|--------|-------|-------|--------|-----------|-------|
| | area | cat* | spp** | area | cat* | spp** | area | cat* | spp** |
| | (ha) | % | % | (ha) | % | % | (ha) | % | % |
| Scots pine | 21 | 1 | 1 | 75 | 2 | 0 | 96 | 2 | 0 |
| Corsican pine | 93 | 4 | 3 | 9 | 0 | 0 | 103 | 2 | 0 |
| Lodgepole pine | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Sitka spruce | 527 | 24 | 18 | 613 | 16 | 3 | 1,140 | 19 | 4 |
| Norway spruce | 177 | 8 | 6 | 66 | 2 | 0 | 243 | 4 | 1 |
| European larch | 31 | 1 | 1 | 298 | 8 | 1 | 329 | 5 | 1 |
| Jap/Hybrid larch | 126 | 6 | 4 | 279 | 7 | 1 | 405 | 7 | 2 |
| Douglas fir | 719 | 33 | 25 | 1,121 | 29 | 5 | 1,840 | 30 | 7 |
| Other conifers | 105 | 5 | 4 | 737 | 19 | 3 | 842 | 14 | 3 |
| Mixed conifers | 407 | 18 | 14 | 657 | 17 | 3 | 1,064 | 18 | 4 |
| Total conifers | 2,207 | 100 | 76 | 3,856 | 100 | 17 | 6,062 | 100 | 24 |
| Oak | 10 | 1 | 0 | 2,376 | 13 | 11 | 2,386 | 12 | 9 |
| Beech | 42 | 6 | 1 | 534 | 3 | 2 | 576 | 3 | 2 |
| Sycamore | 0 | 0 | 0 | 569 | 3 | 3 | 569 | 3 | 2 |
| Ash | 0 | 0 | 0 | 652 | 3 | 3 | 652 | 3 | 3 |
| Birch | 0 | 0 | 0 | 211 | 1 | 1 | 211 | 1 | 1 |
| Poplar | 0 | 0 | 0 | 5 | 0 | 0 | 5 | 0 | 0 |
| Sweet chestnut | 0 | 0 | 0 | 116 | 1 | 1 | 116 | 1 | 0 |
| Elm | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Other broadleaves | 0 | 0 | 0 | 1,364 | 7 | 6 | 1,364 | 7 | 5 |
| Mixed broadleaves | 629 | 92 | 22 | 12,926 | 69 | 57 | 13,556 | 70 | 53 |
| Total broadleaves | 681 | 100 | 24 | 18,752 | 100 | 83 | 19,433 | 100 | 76 |
| Total - all species | 2,888 | | 100 | 22,607 | | 100 | 25,496 | | 100 |
| Felled | 21 | | | 38 | | | 59 | | |
| Total High Forest | 2,909 | | | 22,645 | | | 25,555 | | |

^{*}cat: species percentage of Conifer or Broadleaved in the ownership category **spp: percentage of all species in the ownership category

- In addition to the areas shown there are 656 ha of other areas integral to the woodland not stocked with tree species.
- 2. The standard errors of the all ownerships area estimates for the most common species or species groups are as follows;

| Conifers | 7% |
|-------------------|-----|
| Broadleaves | 3% |
| Sitka spruce | 22% |
| Oak | 14% |
| Other broadleaves | 19% |

- 3. Mixtures: where possible the species in mixtures have been separately recorded. Where this has not been possible they were described as 'Mixed conifers' or 'Mixed broadleaves'.
- 4. Confidence Intervals: where the standard errors of these summary measures are 10% or less, the confidence Intervals will be approximately symmetrical; the true value is expected to be within +/- one standard error for about 68% (or about two-thirds) of all cases, and within +/- two standard errors for about 95% of all cases. Where percentage standard errors are larger, e.g. for less common species or more variable species composition, the confidence intervals will be less symmetrical (and wider).

Area of High Forest by principal species and ownership




Table 9b Area of High Forest by principal species, ownership and category

| Species | Forest | ry Commi | ission | | Other | | All | ownership | os |
|---------------------|-----------|-----------|---------------|-----------|-----------|---------------|-----------|-----------|--------|
| | cat. 1 | cat. 2 | Total (ha) | cat. 1 | cat. 2 | Total (ha) | cat. 1 | cat. 2 | Total |
| Scots pine | 21 | 0 | (na) 21 | 75 | 0 | (na) 75 | 96 | 0 | (ha) |
| Corsican pine | 93 | 0 | 93 | 9 | 0 | 9 | 103 | 0 | 103 |
| Lodgepole pine | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Sitka spruce | 527 | 0 | 527 | 613 | 0 | 613 | 1,140 | 0 | 1,140 |
| Norway spruce | 177 | 0 | 177 | 47 | 19 | 66 | 224 | 19 | 243 |
| European larch | 31 | 0 | 31 | 298 | 0 | 298 | 329 | 0 | 329 |
| Jap/Hybrid larch | 126 | 0 | 126 | 279 | 0 | 279 | 405 | 0 | 405 |
| Douglas fir | 719 | 0 | 719 | 1,121 | 0 | 1,121 | 1,840 | 0 | 1,840 |
| Other conifers | 105 | 0 | 105 | 737 | 0 | 737 | 842 | 0 | 842 |
| Mixed conifers | 407 | 0 | 407 | 593 | 64 | 657 | 1,000 | 64 | 1,064 |
| Total conifers | 2,207 | 0 | 2,207 | 3,773 | 83 | 3,856 | 5,980 | 83 | 6,062 |
| Oak | 0 | 10 | 10 | 1,972 | 404 | 2,376 | 1,972 | 415 | 2,386 |
| Beech | 42 | 0 | 42 | 484 | 50 | 534 | 526 | 50 | 576 |
| Sycamore | 0 | 0 | 0 | 524 | 45 | 569 | 524 | 45 | 569 |
| Ash | 0 | 0 | 0 | 591 | 60 | 652 | 591 | 60 | 652 |
| Birch | 0 | 0 | 0 | 193 | 18 | 211 | 193 | 18 | 211 |
| Poplar | 0 | 0 | 0 | 5 | 0 | 5 | 5 | 0 | 5 |
| Sweet chestnut | 0 | 0 | 0 | 76 | 39 | 116 | 76 | 39 | 116 |
| Elm | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Other broadleaves | 0 | 0 | 0 | 141 | 1,223 | 1,364 | 141 | 1,223 | 1,364 |
| Mixed broadleaves | 262 | 367 | 629 | 9,022 | 3,905 | 12,926 | 9,284 | 4,271 | 13,556 |
| Total broadleaves | 304 | 377 | 681 | 13,007 | 5,744 | 18,752 | 13,311 | 6,122 | 19,433 |
| Total - all species | 2,511 | 377 | 2,888 | 16,780 | 5,827 | 22,607 | 19,291 | 6,205 | 25,496 |


1. The standard errors of the all ownerships area estimates for the most common species or species groups (in all woodland types) are as follows

| | Category 1* Cate | egory 2* | Iotal High Forest | |
|-------------------|------------------|----------|----------------------|------------------------------|
| Conifers | 7% | 53% | 7% | |
| Broadleaves | 4% | 5% | 3% | |
| Sitka spruce | 22% | - | 22% | |
| Oak | 15% | 33% | 14% | *See Glossary for Category 1 |
| Other broadleaves | 47% | 17% | 19% | and Category 2 descriptions |

^{2.} Where the standard errors of these summary measures are 10% or less, the confidence intervals will be approximately symmetrical; the the true value is expected to be within +/- one standard error for about 68% (or about two-thirds) of all cases, and within +/- two standard errors for about 95% of all cases. Where percentage standard errors are larger, e.g. for less common species or more variable species composition, the confidence intervals will be less symmetrical (and wider).

^{3.} Where possible the species in mixtures have been separately recorded. Where this has not been possible they were described as 'Mixed conifers' or 'Mixed broadleaves'.

High Forest Category 1 - Area by principal species and ownership



High Forest Category 2 - Area by principal species and ownership





 Table 10a
 High Forest Category 1 - Area by principal species and planting year class

| Species | | | | | Plo | ınting y | ear cla | ss* | | | | | Total (ha) |
|---------------------|---------------|---------------|---------------|----------------|---------------|----------------|---------------|---------------|----------------|---------------|---------------|--------------|---------------|
| | 1991- 1997 | 1981- 1990 | 1971- 1980 | 1961 - 1970 | 1951- 1960 | 1941 - 1950 | 1931- 1940 | 1921- 1930 | 1911 - 1920 | 1901- 1910 | 1861- 1900 | Pre- 1861 | |
| Scots pine | 9 | O | O | 0 | 10 | 10 | 0 | 4/ | 19 | O | 0 | O | 96 |
| Corsican pine | 9 | 0 | 0 | 93 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 103 |
| Lodgepole pine | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Sitka spruce | 16 | 208 | 87 | 276 | 172 | 381 | 0 | 0 | 0 | 0 | 0 | 0 | 1,140 |
| Norway spruce | 0 | 31 | 0 | 34 | 0 | 106 | 52 | 0 | 0 | 0 | 0 | 0 | 224 |
| European larch | 9 | 14 | 60 | 105 | 72 | 68 | 0 | 0 | 0 | 0 | 0 | 0 | 329 |
| Jap/Hybrid larch | 0 | 13 | 94 | 67 | 201 | 0 | 30 | 0 | 0 | 0 | 0 | 0 | 405 |
| Douglas fir | 288 | 216 | 323 | 298 | 336 | 282 | 96 | 0 | 0 | 0 | 0 | 0 | 1,840 |
| Other conifers | 0 | 60 | 129 | 9 | 226 | 135 | 0 | 52 | 0 | 27 | 68 | 136 | 842 |
| Mixed conifers | 0 | 133 | 303 | 75 | 223 | 110 | 31 | 0 | 19 | 0 | 34 | 72 | 1,000 |
| Total conifers | 332 | 674 | 997 | 959 | 1,240 | 1,092 | 210 | 99 | 38 | 27 | 102 | 209 | 5,980 |
| Oak | 27 | 0 | 41 | 61 | 59 | 175 | 232 | 93 | 451 | 89 | 386 | 355 | 1,972 |
| Beech | 42 | 24 | 0 | 59 | 23 | 12 | 0 | 7 | 57 | 32 | 200 | 71 | 526 |
| Sycamore | 43 | 0 | 19 | 50 | 77 | 39 | 72 | 28 | 36 | 72 | 86 | 0 | 524 |
| Ash | 51 | 0 | 7 | 57 | 36 | 27 | 81 | 3 | 24 | 0 | 264 | 41 | 591 |
| Birch | 0 | 0 | 8 | 73 | 24 | 60 | 0 | 27 | 0 | 0 | 0 | 0 | 193 |
| Poplar | 0 | 0 | 0 | 0 | 0 | 0 | 5 | 0 | 0 | 0 | 0 | 0 | 5 |
| Sweet chestnut | 38 | 0 | 0 | 0 | 0 | 9 | 6 | 0 | 0 | 0 | 4 | 20 | 76 |
| Elm | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Other broadleaves | 51 | 0 | 8 | 37 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 44 | 141 |
| Mixed broadleaves | 216 | 202 | 16 | 36 | 57 | 832 | 570 | 2,320 | 1,831 | 848 | 976 | 1,380 | 9,284 |
| Total broadleaves | 469 | 226 | 100 | 373 | 275 | 1,156 | 966 | 2,478 | 2,399 | 1,042 | 1,917 | 1,911 | 13,311 |
| Total - all species | 801 | 900 | 1,096 | | | | | | | | | | |

^{*}Age determined from records where these were available. Where records were not available or were clearly inaccurate age-class was assigned by reference to similar crops of known age in the locality.

High Forest Category 1 - Area by planting year class



1. Most of the planting year classes cover 10 years, 1991-1997 is 7 years, and the classes prior to 1901 are 40 years or more.


23

 Table 10b
 High Forest Category 1 - Forestry Commission: area by principal species and planting year classes

| Species | | | | | Plo | inting y | ear cla | ss* | | | | | Total (ha) |
|---------------------|---------------|---------------|---------------|----------------|---------------|----------------|---------------|---------------|----------------|---------------|---------------|--------------|---------------|
| | 1991- 1997 | 1981- 1990 | 1971- 1980 | 1961 - 1970 | 1951- 1960 | 1941 - 1950 | 1931- 1940 | 1921- 1930 | 1911 - 1920 | 1901- 1910 | 1861- 1900 | Pre- 1861 | |
| Scots pine | 0 | 0 | 0 | 0 | 10 | 10 | 0 | 0 | 0 | 0 | 0 | 0 | 21 |
| Corsican pine | 0 | 0 | 0 | 93 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 93 |
| Lodgepole pine | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Sitka spruce | 0 | 208 | 73 | 16 | 0 | 231 | 0 | 0 | 0 | 0 | 0 | 0 | 527 |
| Norway spruce | 0 | 31 | 0 | 21 | 0 | 72 | 52 | 0 | 0 | 0 | 0 | 0 | 177 |
| European larch | 0 | 0 | 0 | 0 | 0 | 31 | 0 | 0 | 0 | 0 | 0 | 0 | 31 |
| Jap/Hybrid larch | 0 | 0 | 73 | 10 | 42 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 126 |
| Douglas fir | 168 | 83 | 218 | 31 | 0 | 177 | 42 | 0 | 0 | 0 | 0 | 0 | 719 |
| Other conifers | 0 | 0 | 0 | 0 | 0 | 52 | 0 | 52 | 0 | 0 | 0 | 0 | 105 |
| Mixed conifers | 0 | 0 | 209 | 0 | 104 | 63 | 31 | 0 | 0 | 0 | 0 | 0 | 407 |
| Total conifers | 168 | 322 | 573 | 172 | 156 | 637 | 126 | 52 | 0 | 0 | 0 | 0 | 0 |
| Oak | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Beech | 0 | 0 | 0 | 31 | 0 | 0 | 0 | 0 | 0 | 0 | 10 | 0 | 42 |
| Sycamore | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Ash | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Birch | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Poplar | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Sweet chestnut | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Elm | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Other broadleaves | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Mixed broadleaves | 0 | 21 | 0 | 0 | 21 | 157 | 31 | 31 | 0 | 0 | 0 | 0 | 262 |
| Total broadleaves | 0 | 21 | 0 | 31 | 21 | 157 | 31 | 31 | 0 | 0 | 10 | 0 | 304 |
| Total - all species | 168 | 343 | 573 | 203 | 177 | 795 | 157 | 84 | 0 | 0 | 10 | 0 | 2,511 |

^{*}Age determined from records where these were available. Where records were not available or were clearly inaccurate age-class was assigned by reference to similar crops of known age in the locality.

High Forest Category 1 - Forestry Commission: area by planting year class



1. Most of the planting year classes cover 10 years, 1991-1997 is 7 years, and the classes prior to 1901 are 40 years or more.

Reference Date: 31 March 1997


25

Table 10c High Forest Category 1 - Other ownership: area by principal species and planting year classes

| Species | | | | | Plo | ınting y | ear cla | SS* | | | | | Total (ha) |
|---------------------|---------------|---------------|---------------|----------------|---------------|----------------|---------------|---------------|----------------|---------------|---------------|--------------|---------------|
| | 1991- 1997 | 1981- 1990 | 1971- 1980 | 1961 - 1970 | 1951- 1960 | 1941 - 1950 | 1931- 1940 | 1921- 1930 | 1911 - 1920 | 1901- 1910 | 1861- 1900 | Pre- 1861 | |
| Scots pine | 9 | 0 | 0 | 0 | 0 | 0 | 0 | 47 | 19 | 0 | 0 | 0 | 75 |
| Corsican pine | 9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 9 |
| Lodgepole pine | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Sitka spruce | 16 | 0 | 14 | 260 | 172 | 150 | 0 | 0 | 0 | 0 | 0 | 0 | 613 |
| Norway spruce | 0 | 0 | 0 | 13 | 0 | 34 | 0 | 0 | 0 | 0 | 0 | 0 | 47 |
| European larch | 9 | 14 | 60 | 105 | 72 | 37 | 0 | 0 | 0 | 0 | 0 | 0 | 298 |
| Jap/Hybrid larch | 0 | 13 | 21 | 56 | 159 | 0 | 30 | 0 | 0 | 0 | 0 | 0 | 279 |
| Douglas fir | 120 | 134 | 105 | 267 | 336 | 104 | 55 | 0 | 0 | 0 | 0 | 0 | 1,121 |
| Other conifers | 0 | 60 | 129 | 9 | 226 | 83 | 0 | 0 | 0 | 27 | 68 | 136 | 737 |
| Mixed conifers | 0 | 133 | 94 | 75 | 119 | 47 | 0 | 0 | 19 | 0 | 34 | 72 | 593 |
| Total conifers | 165 | 353 | 423 | 787 | 1,084 | 455 | 85 | 47 | 38 | 27 | 102 | 209 | 3,773 |
| Oak | 27 | 0 | 41 | 61 | 59 | 175 | 232 | 93 | 451 | 89 | 386 | 355 | 1,972 |
| Beech | 42 | 24 | 0 | 27 | 23 | 12 | 0 | 7 | 57 | 32 | 190 | 71 | 484 |
| Sycamore | 43 | 0 | 19 | 50 | 77 | 39 | 72 | 28 | 36 | 72 | 86 | 0 | 524 |
| Ash | 51 | 0 | 7 | 57 | 36 | 27 | 81 | 3 | 24 | 0 | 264 | 41 | 591 |
| Birch | 0 | 0 | 8 | 73 | 24 | 60 | 0 | 27 | 0 | 0 | 0 | 0 | 193 |
| Poplar | 0 | 0 | 0 | 0 | 0 | 0 | 5 | 0 | 0 | 0 | 0 | 0 | 5 |
| Sweet chestnut | 38 | 0 | 0 | 0 | 0 | 9 | 6 | 0 | 0 | 0 | 4 | 20 | 76 |
| Elm | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Other broadleaves | 51 | O | 8 | 3/ | 0 | O | O | U | O | O | O | 44 | 141 |
| Mixed broadleaves | 216 | 181 | 16 | 36 | 36 | 675 | 539 | 2,288 | 1,831 | 848 | 976 | 1,380 | 9,022 |
| Total broadleaves | 469 | 205 | 100 | 341 | 254 | 998 | 935 | 2,446 | 2,399 | 1,042 | 1,907 | 1,911 | 13,007 |
| Total - all species | 634 | 558 | 523 | 1,128 | 1,338 | 1,454 | 1,019 | 2,493 | 2,437 | 1,069 | 2,008 | 2,120 | 16,780 |

^{*}Age determined from records where these were available. Where records were not available or were clearly inaccurate age-class was assigned by reference to similar crops of known age in the locality.

High Forest Category 1 - Other Ownership: area by planting year class



1. Most of the planting year classes cover 10 years, 1991-1997 is 7 years, and the classes prior to 1901 are 40 years or more.

 Table 11 High Forest: principal species by planting year class

| Planting year class | First | % | Second | % | Third | % |
|---------------------|-------------------|----|-------------------|----|-------------------|----|
| 1991-97 | Douglas fir | 34 | Mixed broadleaves | 27 | Sweet chestnut | 8 |
| 1981-90 | Mixed broadleaves | 24 | Douglas fir | 20 | Sitka spruce | 19 |
| 1971-80 | Douglas fir | 28 | Mixed conifer | 26 | Other conifer | 11 |
| 1961-70 | Douglas fir | 18 | Sitka spruce | 16 | Mixed broadleaves | 13 |
| 1951-60 | Douglas fir | 20 | Other conifer | 13 | Mixed conifer | 13 |
| 1941-50 | Mixed broadleaves | 48 | Other broadleaves | 19 | Sitka spruce | 9 |
| 1931-40 | Mixed broadleaves | 68 | Oak | 11 | Douglas fir | 4 |
| 1921-30 | Mixed broadleaves | 90 | Oak | 3 | Other conifers | 2 |
| 1911-20 | Mixed broadleaves | 75 | Oak | 19 | Beech | 2 |
| 1901-10 | Mixed broadleaves | 84 | Oak | 6 | Sycamore | 5 |
| 1861-1900 | Mixed broadleaves | 49 | Oak | 23 | Ash | 10 |
| Pre 1861 | Mixed broadleaves | 65 | Oak | 16 | Other conifers | 6 |
| All years | Mixed broadleaves | 53 | Oak | 9 | Douglas fir | 7 |

^{1.} Principal species as a percentage of area in the planting year class.


Table 12 Ownership type* by area and percentage

| Ownership type | Area (ha) | % |
|------------------------------------|-----------|-------|
| Personal | 14,545 | 54.3 |
| Business | 5,302 | 19.8 |
| Forestry or timber business | 0 | 0.0 |
| Charity | 1,969 | 7.4 |
| Local Authority | 1,022 | 3.8 |
| Other public (not FC) | 852 | 3.2 |
| Forestry Commission | 3,035 | 11.3 |
| Community ownership or common land | 0 | 0.0 |
| Unidentified | 47 | 0.2 |
| Total | 26,771 | 100.0 |

^{*} This table is produced from data contributed on a voluntary basis by owners or their representatives.

Ownership type by area





RESULTS FROM THE SURVEY OF SMALL WOODLAND AND TREES (SSWT)

Survey Method

The land area of England was stratified into coastal and inland 1 km x 1 km squares and a random sample of 1 km² plots were then selected, representing around 1% of the land area. 1:25 000 scale aerial photos were then used to identify features in each sample square. Each 1 km² was then divided into 16 parts, and two of these were selected at random for field data collection. Data was collected on Small Woodlands (0.10 - <2.00 ha), Linear Features, Groups and Individual Trees. The survey did not collect information from areas of developed land of 2 hectares or more.

| Table 13: | Summary of information from the Survey of Small Woodland and Trees |
|-----------|--|
| Table 14: | Woodland area by feature type and woodland size |
| Table 15: | Numbers of live trees outside woodland by species and feature type |
| Table 16: | Numbers of dead trees outside woodland by species and feature type |
| Table 17: | Numbers of live trees outside woodland by species and height band |
| Table 18: | Numbers of Groups by group size |

Note: The figures in many of the tables may not add due to rounding



Table 13 Summary of information from the Survey of Small Woodlands and Trees

| Feature type | Number of features | Total | Unit |
|------------------------|--------------------|-----------|----------------------|
| Small Woods | 164 | 98 | Area (ha) |
| Wide Linear Features | 0 | 0 | Area (ha) |
| Wide Linear Features | 0 | 0 | Length (Km) |
| Narrow Linear Features | 4,900 | 295 | Length (Km) |
| Narrow Linear Features | 4,900 | 186,700 | Number of live trees |
| Groups | 96,400 | 1,022,400 | Number of live trees |
| Individual Trees | 59,600 | 59,600 | Number of live trees |

See Glossary for definitions of feature types.

 Table 14
 Woodland area by feature type and woodland size

| Feature type | Woodland size (ha) | | Total area | Number of | Mean size |
|----------------------|--------------------|-------------|------------|-----------|-----------|
| | 0.1 - <0.25 | 0.25 - <2.0 | (ha) | features | (ha) |
| Small Woods | 5 | 93 | 98 | 164 | 0.60 |
| Wide Linear Features | 0 | 0 | 0 | 0 | 0.00 |
| Total | 5 | 93 | 98 | 164 | 0.60 |

^{1.} See Glossary for definitions of feature types.

Table 15 Numbers of live trees outside woodland by species and feature type (000's trees)

| Species | | Feature | e type | | | Percent of | total trees |
|---------------------|-------------------|--------------|--------|------------------------------|---------------------|------------|-------------|
| | Boundary Trees | Middle Trees | Groups | Narrow Linear Features | Total live trees | Category | Species |
| Pine | 0.8 | 0.0 | 1.6 | 1.9 | 4.3 | 5.1 | 0.3 |
| Spruce | 0.0 | 0.0 | 2.5 | 0.0 | 2.5 | 3.0 | 0.2 |
| Larch | 0.0 | 0.0 | 1.6 | 0.0 | 1.6 | 1.9 | 0.1 |
| Cypress | 0.0 | 0.8 | 13.1 | 0.0 | 13.9 | 16.5 | 1.1 |
| Other conifers | 0.8 | 1.6 | 51.5 | 7.8 | 61.7 | 73.5 | 4.9 |
| Total conifers | 1.6 | 2.5 | 70.3 | 9.7 | 84.0 | 100.0 | 6.6 |
| Oak | 6.5 | 1.6 | 80.9 | 9.7 | 98.7 | 8.3 | 7.8 |
| Beech | 0.8 | 0.0 | 8.2 | 12.6 | 21.6 | 1.8 | 1.7 |
| Sycamore | 8.2 | 0.0 | 263.1 | 32.1 | 303.4 | 25.6 | 23.9 |
| Ash | 11.4 | 4.1 | 139.7 | 4.9 | 160.1 | 13.5 | 12.6 |
| Birch | 1.6 | 1.6 | 7.4 | 0.0 | 10.6 | 0.9 | 0.8 |
| Poplar | 1.6 | 0.0 | 0.0 | 0.0 | 1.6 | 0.1 | 0.1 |
| Sweet chestnut | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Horse chestnut | 0.8 | 0.0 | 0.0 | 0.0 | 0.8 | 0.1 | 0.1 |
| Alder | 0.0 | 0.0 | 8.2 | 0.0 | 8.2 | 0.7 | 0.6 |
| Lime | 0.8 | 0.0 | 0.0 | 0.0 | 0.8 | 0.1 | 0.1 |
| Elm | 0.0 | 0.0 | 67.8 | 0.0 | 67.8 | 5.7 | 5.3 |
| Willow | 0.8 | 0.0 | 113.6 | 75.9 | 190.3 | 16.1 | 15.0 |
| Other broadleaves | 14.7 | 0.8 | 263.1 | 41.8 | 320.4 | 27.1 | 25.3 |
| Total broadleaves | 47.4 | 8.2 | 952.1 | 177.0 | 1184.3 | 100.0 | 93.3 |
| Total - all species | 49.0 | 10.7 | 1022.4 | 186.7 | 1268.7 | | 100.0 |

Percentages

Category: species percentage of conifer or broadleaved. Species: percentage of all species.

2. The standard errors of the total tree number estimates for these feature types are:

Individual Trees24%Groups26%Narrow Linear Features58%

3. See Glossary tor definitions of teature types.

 Table 16
 Numbers of dead trees outside woodland by species and feature type (000's trees)

| | | Featur | e type | | | Percent c | of total trees |
|---------------------|-------------------|--------------|--------|------------------------------|---------------------|-----------|----------------|
| Species | Boundary Trees | Middle Trees | Groups | Narrow Linear Features | Total dead trees | Category | Species |
| Pine | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Spruce | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Larch | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Cypress | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Other conifers | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Total conifers | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Oak | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Beech | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Sycamore | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Ash | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Birch | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Poplar | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Sweet chestnut | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Horse chestnut | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Alder | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Lime | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Elm | 0.8 | 0.0 | 0.0 | 0.0 | 0.8 | 100.0 | 100.0 |
| Willow | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Other broadleaves | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Total broadleaves | 0.8 | 0.0 | 0.0 | 0.0 | 0.8 | 100.0 | 100.0 |
| Total - all species | 0.8 | 0.0 | 0.0 | 0.0 | 0.8 | | 100.0 |

^{1.} See Glossary for definitions of feature types.

Table 17 Numbers of live trees outside woodland by species and height band (000's trees)

| Species | | Total live trees | | | |
|---------------------|-------|------------------|-------|-----|---------|
| | 2-5 | 5-15 | 15-20 | >20 | |
| Pine | 0.8 | 2.8 | 0.8 | 0.0 | 4.4 |
| Spruce | 2.5 | 0.0 | 0.0 | 0.0 | 2.5 |
| Larch | 0.0 | 1.6 | 0.0 | 0.0 | 1.6 |
| Cypress | 0.0 | 13.9 | 0.0 | 0.0 | 13.9 |
| Other conifers | 1.6 | 60.1 | 0.0 | 0.0 | 61.7 |
| Total conifers | 4.9 | 78.4 | 0.8 | 0.0 | 84.1 |
| Oak | 20.4 | 56.5 | 21.8 | 0.0 | 98.7 |
| Beech | 0.0 | 2.5 | 19.2 | 0.0 | 21.7 |
| Sycamore | 0.8 | 288.7 | 13.9 | 0.0 | 303.4 |
| Ash | 0.8 | 138.1 | 21.2 | 0.0 | 160.1 |
| Birch | 0.8 | 4.9 | 4.9 | 0.0 | 10.6 |
| Poplar | 0.0 | 1.6 | 0.0 | 0.0 | 1.6 |
| Sweet chestnut | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Horse chestnut | 0.0 | 0.8 | 0.0 | 0.0 | 0.8 |
| Alder | 0.0 | 8.2 | 0.0 | 0.0 | 8.2 |
| Lime | 0.0 | 0.0 | 0.8 | 0.0 | 0.8 |
| Elm | 3.3 | 64.6 | 0.0 | 0.0 | 67.9 |
| Willow | 0.0 | 190.3 | 0.0 | 0.0 | 190.3 |
| Other broadleaves | 96.1 | 224.4 | 0.0 | 0.0 | 320.5 |
| Total broadleaves | 122.2 | 980.6 | 81.8 | 0.0 | 1,184.6 |
| Total - all species | 127.1 | 1,059.0 | 82.6 | 0.0 | 1,268.7 |

Table 18 Number of Groups by group size

| Number of trees per Group* | Number of Groups (000's) |
|-------------------------------|-----------------------------|
| 2 | 10 |
| 3-5 | 20 |
| 6-10 | 26 |
| 11-20 | 20 |
| 21-50 | 16 |
| 51-100 | 5 |
| >100 | 0 |
| Total | 96 |

^{*}The size of the group is determined by the total number of trees, live plus dead.

COMPARISON OF RESULTS WITH THE 1980 CENSUS AND PREVIOUS SURVEYS

Survey Method

The 1980 Census and 1997 Inventory were undertaken using very different sampling methods.

Inventory practice and technology have moved on since the 1980 Census; this has led to changes in sampling methodology, scope and woodland definitions. For example, the Main Woodland Survey used the digital woodland map, created from aerial photos as a basis for sampling whereas the 1980 Census relied only on the woodland shown on the 1:50,000 Ordnance Survey map. Also in contrast to the 1980 Census, the Survey of Small Woodland and Trees did not record information within developed land e.g. residential or industrial areas of 2 or more hectares.

Where possible adjustments have been made to both the 1980 Census and the Inventory to achieve the nearest available comparison. The apparent changes indicated in the following tables and charts should therefore be treated with caution, particularly where areas are small.

Table 19: Comparison of woodland area

between 1980 Census and 1997 Inventory

Table 20: Comparison of High Forest area by species

between 1980 Census and 1997 Inventory

Chart: Comparison of High Forest area by species

between 1980 Census and 1997 Inventory

Comparison of High Forest Category 1 area by planting year class Table 21:

between 1980 Census and 1997 Inventory

Comparison of High Forest Category 1 area by planting year class Chart:

between 1980 Census and 1997 Inventory

Table 22: Comparison of numbers of live trees outside woodland

between 1980 Census and 1997 Inventory

Table 23: Comparison of density of non-woodland features

between 1980 Census and 1997 Inventory

Woodland cover

Chart Change in woodland cover through time (1890 – 2000)

Maps: Woodland by county through time (1895 – 1998)

Note: The figures in many of the tables may not add due to rounding



Table 19 Comparison of woodland area between 1980 Census and 1997 Inventory

| Woodland size (ha) | 1980 Census woodland area | | 1997 In woodla | Change (%) | |
|-----------------------|------------------------------|------|-------------------|---------------|-----|
| | (ha) | (%) | (ha) | (%) | (%) |
| 2.0 or more | 16,297 | 89.0 | 26,771 | 99.7 | 64 |
| 0.25 - <2.0 | 2,004 | 11.0 | 93 | 0.3 | -95 |
| Total | 18,301 | | 26,864 | | 47 |
| % Woodland land cover | 5.1 | | 7.5 | | |

- 1. Differences in sampling methodology may account for some of the apparent differences.
- The above figures from the 1997 Inventory exclude woodland between 0.1 and <0.25 ha, thereby matching the scope of the 1980 Census.
 The 1997 figures above will therefore not match those in the previous sections of the report.
- 3. Land area used to calculate woodland cover percent (1997), 356,466 ha, was based on the 1991 Census of Population digital boundaries.
- Land area used to calculate woodland cover percent (1980), 356,422 ha,
 (Ordnance Survey data)

41

Table 20 Comparison of High Forest area by species between 1980 Census and 1997 Inventory

| Species | 1980 Census woodland area (ha) | 1997 Inventory woodland area (ha) | Change (%) |
|-------------------|-----------------------------------|--------------------------------------|---------------|
| Scots pine | 115 | 97 | -16 |
| Corsican pine | 46 | 103 | 125 |
| Lodgepole pine | 220 | 0 | -100 |
| Sitka spruce | 1,585 | 1,140 | -28 |
| Norway spuce | 338 | 243 | -28 |
| European larch | 82 | 329 | 301 |
| Jap/Hybrid larch | 909 | 413 | -55 |
| Douglas fir | 1,127 | 1,840 | 63 |
| Other conifers | 493 | 846 | 72 |
| Mixed conifers | 275 | 1,064 | 287 |
| Total conifers | 5,190 | 6,075 | 17 |
| Oak | 5,941 | 2,395 | -60 |
| Beech | 1,363 | 606 | -56 |
| Sycamore | 783 | 582 | -26 |
| Ash | 1,099 | 659 | -40 |
| Birch | 173 | 211 | 22 |
| Poplar | 26 | 5 | -81 |
| Sweet chestnut | 88 | 116 | 32 |
| Elm | 1 | 4 | 310 |
| Other broadleaves | 1,400 | 1,379 | -1 |
| Mixed broadleaves | 768 | 13,558 | 1666 |
| Total broadleaves | 11,642 | 19,515 | 68 |
| Total all species | 16,832 | 25,590 | 52 |
| Felled | 516 | 59 | -89 |
| Total High Forest | 17,348 | 25,649 | 48 |

^{1.} Differences in sampling methodology may account for some of the apparent differences.

^{2.} In the 1980 Census the areas assigned to species included any associated open space such as roads and rides. In the Inventory open spaces are separately identified and the overall proportion is 2.4% (Table 2). To obtain meaningful comparisons between the two datasets the 1980 Census data have therefore been reduced by 2.4%.

The above figures from the 1997 Inventory exclude woodland between 0.1 and <0.25 ha, thereby matching the scope of the 1980 Census.
 The 1997 figures above will therefore not match those in the previous sections of the report.

^{4.} The 1980 figures include scrub to enable comparison

Comparison of High Forest area by species between 1980 Census and 1997 Inventory




Table 21 Comparison of High Forest Category 1 area by planting year class between 1980 Census and 1997 Inventory

| Planting year class | 1980 Census woodland area (ha) | 1997 Inventory woodland area (ha) | Change (%) |
|---------------------|-----------------------------------|--------------------------------------|---------------|
| 1991-1997 | 0 | 839 | see note |
| 1981-1990 | 0 | 900 | see note |
| 1971-1980 | 1,284 | 1,101 | -14 |
| 1961-1970 | 2,249 | 1,332 | -41 |
| 1951-1960 | 1,586 | 1,520 | -4 |
| 1941-1950 | 737 | 2,248 | 205 |
| 1931-1940 | 1,124 | 1,176 | 5 |
| 1921-1930 | 656 | 2,577 | 293 |
| 1911-1920 | 601 | 2,440 | 306 |
| 1901-1910 | 476 | 1,069 | 125 |
| 1861-1900 | 2,227 | 2,030 | -9 |
| Pre 1861 | 770 | 2,120 | 175 |
| Total all years | 11,709 | 19,352 | 65 |

^{1.} The first two classes, 1991-1997 and 1981-1990, cover the period since the 1980 Census and no comparison is therefore available.

^{2.} The definition of High Forest Category 1 in the Inventory does not fully coincide with High Forest as defined in the 1980 Census.

Comparison of High Forest Category 1 area by planting year class between 1980 Census and 1997 Inventory




Table 22 Comparison of numbers of live trees outside woodland between 1980 Census and 1997 Inventory (000's)

| Feature type | 1980 Census | 1997 Inventory | Change (%) |
|------------------------|-------------|----------------|------------|
| Boundary Tree | 148 | 43 | -71 |
| Middle Tree | 81 | 11 | -86 |
| Total Individual Trees | 228 | 55 | -76 |
| Groups | 645 | 780 | 21 |
| Linear Features | 570 | 151 | -74 |
| Total | 1,443 | 985 | -32 |

- The Survey of Small Woodland and Trees did not record information referring to tree
 features (I.e. Individual trees, Groups and Narrow Linear Features) within developed
 land.
- In the 1980 Census hazel, hawthorn, blackthorn and goat willow were excluded, the 1997 Inventory figures have been adjusted accordingly.
 The 1997 figures above will therefore not match those in the previous sections of the report.
- 3. Changes stated in this table are indicative only. Even with adjustments to the 1997 Inventory, the two surveys are not directly comparable 1980 used 7cm diameter at breast height, and 1997 used 2m height, as minimum criteria for inclusion.
- 4. See Glossary for definitions of feature type.

Table 23 Comparison of density of non-woodland features between 1980 Census and 1997 Inventory

| Feature type | 1980 Census | 1997 Inventory | Change (%) |
|-------------------------------|-------------|----------------|------------|
| Individual Trees (per sq km) | 64.0 | 15.3 | -76 |
| Groups (per sq km) | 40.0 | 22.0 | -45 |
| Linear Features (m per sq km) | 297.4 | 82.7 | -72 |


- The Survey of Small Woodland and Trees did not record information referring to tree features (I.e. Individual trees, Groups and Narrow Linear Features) within developed land.
- In the 1980 Census hazel, hawthorn, blackthorn and goat willow were excluded, the 1997 Inventory figures have been adjusted accordingly.
 The 1997 figures above will therefore not match those in the previous sections of the report.
- Changes stated in this table are indicative only. Even with adjustments to the 1997 Inventory, the two surveys are not directly comparable - 1980 used 7cm diameter at breast height, and 1997 used 2m height, as minimum criteria for inclusion.
- 4. See Glossary for definitions of feature type.

WOODLAND COVER


Woodland area data is available from Ministry of Agriculture surveys since 1871, and from Forestry Commission national woodland inventories since 1924. The following chart and maps show the changes in woodland area through time.


The maps use the old County structure data of England, as reported on in 1895 and 1947. The data from these counties could not be re-worked for different geographic areas. In contrast, the digital woodland map, which forms the basis of the current inventory, can be analysed for any geographic area.


Change in county woodland cover through time (1890 – 2000)



Map 5 Woodland Cover in England by County through time (1895–1998)







GLOSSARY

Woodland

In the United Kingdom woodland is defined as land with a minimum area of 0.1 ha under stands of trees with, or the potential to achieve, tree crown cover of more than 20%. Areas of open space integral to the woodland are also included. Orchards and urban woodland between 0.1 and 2 ha are excluded. Intervening land-classes such as roads, rivers or pipelines are disregarded if less than 50m in extent. 'Scrubby' vegetation is not Included as a separate category but as Conifer, Broadleaved or Mixed tree types. There is additional information on the quality of woodland within the inventory database.

Woodland of 2 ha and over, and with a minimum width of 50m, is included in the Main Woodland Survey; other woodland and trees are assessed in the Survey of Small Woodland and Trees.

Interpreted Forest Types

The woodland map derived from aerial photographs is differentiated into Interpreted Forest Types (IFTs) which are: Conifer, Broadleaved, Mixed, Coppice, Coppice-with-Standards, Shrubs, Young Trees, Ground Prepared for Planting and Felled. Note that forest types (see below) based on ground survey data are used for reporting purposes because they are more reliable.

High Forest

All woodland except stands managed as Coppice or Coppice-with-Standards with, or with the potential to achieve a tree cover of more than 20%. Two categories of High Forest are recognised:

High Forest Category 1

Stands which are, or could become, capable of producing wood of a size and quality suitable for sawlogs.

High Forest Category 2

Stands of lower quality than High Forest Category 1.

Mixtures

Where possible the species in mixtures have been separately recorded. Where this has not been possible they were described as 'Mixed conifers' or 'Mixed broadleaves'.

Forest Types

Conifer

Woodland containing more than 80% by area of coniferous species.

Broadleaved

Woodland containing more than 80% by area of broadleaved species.

Mixed

A combination of broadleaved and coniferous species where each category occupies at least 20% of the canopy (see note on mixtures above.)

Coppice

Crops of marketable broadleaved species that have at least 2 stems per stool and are either being worked or are capable of being worked on rotation. With the exception of hazel coppice more than half the stems should be capable of producing 1m timber lengths of good form.

Coppice with Standards

Two-storey stands where the overstorey consists of at least 25 stems per ha that are older than the understorey of worked coppice by at least one coppice rotation.

Felled

Woodland areas that have been felled or stands where the stocking has been reduced to less than 20% and where it is expected that these areas will be replanted.

Windblow

Areas of blown woodland which remain uncleared and not regenerated.

Open Space

Areas within a woodland that are not covered by trees but are integral to the woodland such as open areas, streamsides, deer glades, rides and forest roads.

Ownership types

Other Ownership

Woodland other than that owned by, or leased to, the Forestry Commission

- Personal

types of private occupation, e.g. individuals, private family trusts and family partnerships.

- Private forestry or timber business

owned by wood processing industry. This category does not include forest management companies.

- Other private business

occupiers, e.g. companies, partnerships, syndicates and pension funds.

- Local Authority

Region, County, District or other Council

- Other public bodies (not FC)

Government department/agency, nationalised industry, etc.

- Charitable organisations

organisations funded by voluntary public subscription, e.g. National Trust, churches and colleges.

- Community ownership or common land

the common property of all members of the community.

• Forestry Commission

Land owned by or land leased to the Forestry Commission

Feature types

Small Wood

A woodland with an area of 0.1 ha or over but less than 2 ha.

Group

A group containing two or more trees with an area less than 0.1ha.

• Individual Tree

A tree the crown of which has no contact with any other tree crown and which is at least 2m tall. Two types of individual tree are recognised:

- Boundary Tree (an Individual Tree on any boundary)
- Middle Tree (an Individual Tree not on a boundary)

• Linear Feature

A feature with a length of 25 m or more, and one which is at least four times as long as it is broad. It can be up to 50m wide or as narrow as a single line of trees. Two types of Linear Features are recognised:

- Narrow Linear Features (with a width of 16 m or less)
- Wide Linear Features (with a width greater than 16 m)

NOTES





231 Corstorphine Road Edinburgh EH12 7AT

www.forestry.gov.uk