

# NFI woodland ecological condition in England: classification results

National Forest Inventory

**Issued by:** National Forest Inventory, Forestry Commission,  
231 Corstorphine Road, Edinburgh, EH12 7AT

**Date:** Feb 2020

**Enquiries:** Ben Ditchburn, 0300 067 5561  
NFI@forestresearch.gov.uk

**Statistician:** David Ross  
David.Ross@forestresearch.gov.uk

**Website:** [www.forestresearch.gov.uk/inventory](http://www.forestresearch.gov.uk/inventory)  
[www.forestresearch.gov.uk/forecast](http://www.forestresearch.gov.uk/forecast)

## Summary

The National Forest Inventory (NFI) for Great Britain provides information concerning the size, distribution and composition of forests and woodlands including woodland ecological condition. Data enabling the calculation of 15 ecological condition indicators were measured as part of the NFI survey cycle 2010-2015 and compared to a benchmark of a stand of ancient semi-natural woodland (ASNW) in good condition. This enabled woodland stands to be classified as favourable, intermediate or unfavourable in terms of their ecological condition. These results can be calculated by woodland type, habitat type, country and region.

This report contains the classification results broken down by woodland type (native, non-native and 'near-native and fragments'). A series of complementary reports has been produced to describe the methodology used, to set out the underpinning statistics on the condition indicators and classes, and to summarise the results. Figure 1.1 illustrates the report and data products available.

For a brief summary of the work or for full details of the methodology please refer to:

- *NFI Woodland ecological condition in Great Britain: Executive Summary*
- *NFI Woodland ecological condition in Great Britain: Methodology*

For the statistical results for other countries please refer to:

- *NFI Woodland ecological condition in [country<sup>1</sup>]: Statistics*

For the classification results for the individual British countries please refer to:

- *NFI Woodland ecological condition in [country<sup>1</sup>]: Classification Results*

Results for individual priority woodland habitat types can be found in the supporting data:

- *NFI Woodland Ecological Condition in [country]: Supporting Data<sup>2</sup>*


---

<sup>1</sup> There are four separate reports; Great Britain, England, Scotland and Wales

<sup>2</sup> Supporting/additional data is available as MS-Excel® spreadsheets

# NFI woodland condition classifications

**Figure 1.1.** A schematic diagram to illustrate the link between all the reports published by the NFI on the study of woodland ecological condition in Great Britain.



Notes: [green boxes] = published reports; [grey box] = detail available in supporting spreadsheets; [pink box] = over-arching theme. GB = Great Britain.

Ten reports have been published in relation to woodland ecological condition, namely; executive summary, methodology, statistics and classification results:

1. The Executive Summary spans all three topics presenting an overview of the methodology, key results and sign-posting to more detail;
2. The methodology describes the survey methodology and the calculation of the condition scores;
3. The statistics reports describe the key statistical results, one for each of the three countries and Great Britain, broken by native woodland type<sup>1</sup> and;
4. The classification results describe woodland ecological condition (as calculated by the NFI Condition Calculator, see Methodology) one for each of the three countries and Great Britain by woodland type.

<sup>1</sup> The written reports cover native woodland type, the supplementary data includes regional breakdowns and statistics and classification by priority habitat types as well.

## Key findings:

The primary purpose of this report and the supporting data is to provide classifications of woodland ecological condition in native woodland habitat. For comparative purposes classifications for non-native woodland and near native woodland have been supplied.

- The total area of native woodland in Great Britain is estimated at around 1.51 million hectares, which is circa 500 thousand hectares or circa 50% higher than reported in previous estimates (see page 15, Table 3.2).
- The total area of native woodland in England is estimated at around 914 thousand hectares, which is circa 250 thousand hectares or 27% higher than reported in previous 2008 HAP estimates.
- 83 thousand hectares or 9% of native woodland area in England is in favourable condition<sup>1</sup>.
- 815 thousand hectares or 90% of native woodland area in England is in intermediate condition<sup>2</sup>.
- 5 thousand hectares or 0.5% of woodland area in England is in unfavourable condition.
- The principal reason for woodlands falling into unfavourable or intermediate condition is due to the fragmentation of woods, low levels of older trees and veteran trees. For example 99% of native woodland area in England is in unfavourable condition for the presence of veteran trees.
- Other factors have added to the number of stands classified as unfavourable or intermediate condition in England:
  - Herbivore damage is found in many stands, with 38% of woodland area in unfavourable condition for this factor.
  - Deadwood levels are unfavourable for 79% of native woodland area.
  - 10% of native woodland area has unfavourable status for invasive species.
  - 2% of native woodland area has unfavourable status for pests and diseases.
- The regeneration of (native and non-native) young trees in native woodland area in England is positive with 81% as intermediate for regeneration (stand-level) and 19% favourable.
- 68% of native woodland area in England is in favourable condition for the number of native species within their canopy.
- 59% of native woodland area in England is in favourable condition for vertical canopy structure.
- 86% of native woodland area in England is in favourable condition for 'nativeness' of canopy.

---

<sup>1</sup> Section 11, Table 11.1 summarises the WEC indicators and thresholds used to classify stands as favourable, unfavourable and intermediate.

<sup>2</sup> With only one reference point of NFI woodland ecology measurements taken currently, no assessment of condition trends, whether declining or improving, can be made. Future reports utilising the second NFI cycle will address this.

## Contents

1	Introduction.....	7
1.1	Aim .....	7
1.2	The importance of understanding the ecological condition of England’s woodlands .....	7
1.3	Measuring ecological condition .....	8
1.3.1	The NFI Map: an overview	9
1.3.2	NFI woodland sample evaluation: an overview	9
1.3.3	Ecological condition classification	10
1.3.4	Extrapolating NFI field survey statistics to a reporting area	10
1.3.5	The NFI Condition Calculator: an overview	11
1.3.6	Woodland and woodland type definitions	11
2	Results .....	13
2.1	Standard error .....	13
3	Woodland habitat type and woodland native type.....	14
4	Stand-level condition classification results for native woodland area in England .	16
5	Stand-level condition classification results for near native woodland and fragments in England .....	18
6	Stand-level condition classification results for non-native woodland stands in England .....	20
7	Condition scoring distribution.....	22
8	Discussion .....	23
8.1	Woodland habitat area.....	23
8.2	Classification of woodland condition .....	24
8.3	Conclusion.....	26
9	References .....	27
10	Glossary.....	27
11	Annex.....	32
12	NFI national reports .....	34

## Figures

<b>Figure 1.1.</b> A schematic diagram to illustrate the link between all the reports published by the NFI on the study of woodland ecological condition in Great Britain. ....	3
<b>Figure 4.1</b> The proportion of each woodland ecological condition class, for each WEC indicators in native woodland stands in England .....	17
<b>Figure 5.1</b> The proportion of each woodland ecological condition class, for each WEC indicator type in near native woodland and fragments in England .....	19
<b>Figure 6.1</b> The proportion of each woodland ecological condition class, for each WEC indicator type in non-native woodland stands in England.....	21
<b>Figure 7.1</b> The overall distribution of ecological condition class by woodland type in England .....	22

## Tables

<b>Table 3.1</b> Area of woodland by habitat type for each country and region of GB .....	14
<b>Table 3.2</b> Area of woodland by woodland type for each country and region of GB.....	15
<b>Table 4.1</b> Area of native woodland stands in England classified as favourable, intermediate or unfavourable for each of the 15 WEC indicators .....	16
<b>Table 5.1</b> Area of near native woodland and fragments in England classified by woodland ecological condition for each of the 15 WEC indicators .....	18
<b>Table 6.1</b> Area of non-native woodland stands in England classified by woodland ecological condition classification for each of the 15 WEC indicators .....	20
<b>Table 11.1</b> Classification threshold summary .....	32

## 1 Introduction

British woodlands are dynamic entities and their ecological condition changes in response to factors such as changing woodland management, general land use practices and climate change. There is a growing requirement across government, non-governmental organisations and the private sector for a better understanding of the ecological condition of British woodlands, which can be used to inform the targeting of resources and woodland management in support of biodiversity and ecological resilience.

This is the largest and most in-depth direct field-based assessment of woodland ecological condition to have been carried out in England. The study has been run in combination with the [National Forest Inventory](#) (NFI) that records the size, distribution and composition of woodland in Great Britain.

### 1.1 Aim

The purpose of this report is to present the area and proportion of woodland in three woodland ecological condition classes (favourable, intermediate, unfavourable) for native, non-native and 'near-native and fragments' woodland in England. Classification results by woodland habitat type are available in the supporting Excel tables.

### 1.2 The importance of understanding the ecological condition of England's woodlands

There are several drivers for monitoring woodland ecological condition. The primary objective is to provide government with statistics that reflect woodland ecological condition and the main factors that impact condition, so that they can make informed decisions concerning the current state of woodland ecological condition and its management. Secondly, it is the UK's national and international requirement to monitor woodland condition. The UK government signed the following global and pan-European (EU) agreements in 1992 which led to commitments concerning the protection of biodiversity:

- The global Convention on Biological Diversity (CBD; <https://www.cbd.int>). Contracting parties are required to develop and enforce national strategies to identify, conserve and protect existing biodiversity. Article 7 of the convention focuses on the requirement to monitor biodiversity.
- The EU Habitats Directive (Directive 92/43/EEC) aims to promote the maintenance of biodiversity by requiring Member States to take measures to maintain or restore natural habitats and wild species listed on its Annexes to a favourable conservation status (JNCC, 2018). Article 17 of the directive specifically requires members to report an assessment of the conservation status of species and habitats listed on the Annexes of the Directive every 6 years.

## 1.3 Measuring ecological condition

The [National Forest Inventory](#) (NFI) field survey assesses a large, stratified-random sample of woodlands across GB on a 5-year rolling cycle using a standardised protocol. Detailed data on various attributes are collected from approximately 15,100 one-hectare sample squares that are partially or entirely covered by forest, including clear-felled areas, according to the woodland map. The first cycle ran from 2010 to 2015 and the second cycle commenced in 2015 (to be completed in 2020). The survey provides an extensive, in-depth and spatially explicit record of our forests and woodlands.

This report provides a brief overview of the methodology used to assess woodland condition using the first cycle of NFI survey data, but full details are provided in the complementary report, 'NFI woodland ecological condition in Great Britain: Methodology'.

To assess the ecological condition of the NFI survey woodlands, 15 woodland ecological condition (WEC) indicators were devised, reviewed and agreed by the NFI WEC working group, a group of specialists from Natural England, Forestry Commission, Forestry England, Scottish Forestry, Scottish Natural Heritage, Natural Resources Wales and the Welsh Government:

1. Age distribution of trees
2. Herbivore damage
3. Invasive plant species
4. Number of native trees
5. Occupancy of native trees
6. Open space
7. Proportion of favourable land cover
8. Woodland regeneration (stand-level) <sup>1</sup>
9. Woodland regeneration (square-level)
10. Tree health
11. Vegetation and ground flora
12. Woodland vertical structure
13. Veteran trees
14. Volume of deadwood
15. Total area of woodland

---

<sup>1</sup> Component group = Homogeneous areas that are too small (<0.05 ha) to practically map as a discrete section using Geographic Information System (GIS) software in the field, but with most of the same defining characteristics as a section. Section = within each sample square, the forest was stratified into different woodland stands or 'sections'.



# NFI woodland condition classifications

---

Alongside the WEC indicators listed above (i.e. stand-level indicators), total woodland area and woodland area loss were designated as population-level indicators (i.e. national-level indicators) and are reported upon and evaluated separately (see [Forestry Statistics](#)) to the condition classification results presented here. The national-level estimates were derived from the NFI woodland map, augmented by the NFI fieldwork. Both the national-level indicators and stand-level indicators are pertinent to the overall, national picture of habitat condition and should be considered together.

## 1.3.1 The NFI Map: an overview

Data on the location and extent of all forests and woodlands in the UK ( $\geq 0.5$  hectares) is created, stored and maintained by the National Forest Inventory (NFI) as a [digital woodland map](#). The map provides current and historical information on the distribution, type and size of forests and woodland in the UK. The digital woodland map is produced using aerial photography, satellite imagery and administrative records and updated annually.

## 1.3.2 NFI woodland sample evaluation: an overview

NFI sample squares were located within the NFI mapped area using a stratified, random approach. Surveyors classify the area within each one hectare NFI sample square into forest and non-forest. The forested area(s) within the sample square are then further stratified into smaller units of homogenous canopy type based on differences in features such as woodland habitat and tree species, or more subtle factors such as condition and thinning history. Such contiguous 'units' of woodland are referred to as 'stands' for the purposes of these reports. Within each forest stand, information on species, age, management regime and other data is collected to enable assessment of each of the 15 ecological indicators described above.

Typically, there are multiple stands which may also be further divisible within each sample square. This means, from the circa 6,500 sample squares covered by the NFI in England, 17,000 forest stands have been assessed for ecological condition. Within each stand two or three 100 m<sup>2</sup> (0.01 hectare) circular plots were randomly located. Within each of these circular plots, stocking was assessed and species, age, grid location and diameter at breast height (DBH) was recorded for all trees  $\geq 4$  cm DBH. A total of 330,000 trees were measured in England. Transects were also conducted within each circular plot to assess the volume of lying deadwood, seedlings and saplings.

Each of the sample squares are marked on the ground with metal pegs and their exact location recorded by GPS to allow for quality assurance checks and future assessment. All field measurements and information are subject to quality assurance checks.

# NFI woodland condition classifications

---

## 1.3.3 Ecological condition classification

The NFI WEC working group established a process for using the WEC indicator data to classify and score woodlands according to their expected condition. Briefly, the process developed involves the following steps:

1. Collect data on the WEC indicators as part of the NFI field survey.
2. Supply statistics on these indicators e.g. 'x % of woodland stands showed evidence of regeneration'.
3. Using ancient semi-natural woodland (ASNW) in good condition as a benchmark (see below), define thresholds for classifying woodland stands into 'favourable', 'intermediate' or 'unfavourable' status for each WEC indicator.
4. Assign numerical scores to these categories and combine these scores for all WEC indicators to provide an overall condition status score for each woodland stand.
5. Define thresholds to apply to the combined scores in order to classify woodland stands into overall 'favourable', 'intermediate' or 'unfavourable' status.
6. Supply information on the classification and scores of woodlands e.g. 'x % of woodland stands were classified as being in favourable condition status for the regeneration indicator'.
7. Use the results from the first survey cycle as a baseline against which changes in condition can be measured for monitoring purposes using data from future survey cycles.

The indicators and classification thresholds were selected with reference to other work (e.g. the Common Standards Monitoring approach for protected sites (JNCC, 2003)), the best available scientific evidence, expert opinion and each country's current policy needs and targets. For example, to determine a 'favourable' level of seedlings, saplings and small trees, a combination of published evidence and expert opinion was used. Details of thresholds used can be found in the methodology report.

## 1.3.4 Extrapolating NFI field survey statistics to a reporting area

The data collected within each survey sample square (used for the 15 WEC indicators), the derived scores and the classifications, were extrapolated and aggregated to the areas of woodland recorded in the NFI map (e.g. woodland type) using standard statistical survey methodology. The classifications of each stand can also be extrapolated up to any sub class of woodland area, such as woodland type or habitat type (minimum size 30,000 ha) and in turn broken down by any geographic area, such as by country or NFI region, as in this report. This report presents the results for different woodland types (native, near-native and fragments', non-native) for GB, results specifically for GB, Scotland and Wales are presented in companion reports (Figure 1.1). Results for individual priority woodland habitats are available in the supplementary data.

# NFI woodland condition classifications

---

## 1.3.5 The NFI Condition Calculator: an overview

To report on condition using the NFI data, an analytical tool was developed, referred to herein as the NFI 'Condition Calculator'. This tool allows the detailed data recorded in each NFI survey square to be analysed alongside the NFI woodland map and other data. It automatically produces the stand-level condition results per woodland type and aggregated statistics for the reporting area. The advantages of establishing an automated reporting tool are that results can be generated on demand using a consistent approach. The Condition Calculator will therefore allow the data from future cycles of the NFI to be analysed using the same procedures, enabling reliable comparisons for reporting on change.

## 1.3.6 Woodland and woodland type definitions

### **Woodland**

An area of land over 0.5 hectares in extent, with 20% or more tree canopy cover, or the potential to achieve that based upon established trees.

### **Native woodland**

Stands with 50% or more native tree species<sup>1</sup> occupancy in the upper canopy that either:

- Form a discrete woodland parcel with a minimum area of 0.5 ha.
- Form a woodland stand with a minimum area of 0.1 ha that is part of a woodland that is 0.5 ha or larger.

### **Non-native woodland**

Stands with less than 40% native species occupancy sitting within a woodland of any size.

### **Near native and fragments**

Stands that fail to meet the criteria for native or non-native woodland specified above are classified as 'near native and fragments'. Defining this category allows all woodland area to be assessed and reported on for its ecological condition status. Pinpointing these areas of woodland may help inform targeted restoration, as they may represent previously native woodland area that has been overplanted with non-natives.

The near native and fragments woodland type can be subdivided into two subclasses:

1. **Near native:** have a native canopy cover of somewhere between 40% to 49% and thus are 'nearly' native.
2. **Fragments:** have 50% or more native tree species occupancy in the upper canopy but fall under the minimum size threshold of 0.1 ha, falling in the size range 0.05 ha to 0.099 ha.

---

<sup>1</sup> For a list of native species refer to the methodology report.

## NFI woodland condition classifications

---

### **Not determinable**

Areas classified as 'not determinable' apply to woodland areas that cannot be classified due to insufficient tree or other attribute information, such as areas without canopy cover and clear-fell sites with a weak vegetation layer. These form less than 0.5% of the whole woodland population.

## 2 Results

This report gives the scores of the woodland ecological condition assessment in England by woodland type. Given the large volume of data gathered in this study it was decided that the reports would focus on results by country and woodland type. Results by habitat type and region are available in the supporting data Excel spreadsheets.

The nominal baseline date for estimates is an average of 2013 (mid-point of the survey) and the time period over which the full series of field samples were collected was January 2010 to January 2016.

All areas in the tables are rounded to the nearest 100 hectares. In some breakdowns, the estimates in the body of the table may not sum to the quoted total because each individual value, including the total, has been independently generated by the estimation procedure used for results from the NFI sample survey.

### 2.1 Standard error

Along with these estimates, associated sampling standard errors have also been calculated and reported, giving a measure of accuracy, conditional upon the underlying assumptions. The sampling standard error will account for random variation arising from the selection of the sample and random measurement errors. However, standard error will not account for any systematic biases in the field measurements, but it is unlikely that any substantial biases of this nature are present in the survey data because the quality assurance processes account for this. Any standard error greater than 25% is reported in amber text and represents a lower degree of assurance in the estimates.

## 3 Woodland habitat type and woodland native type

This is the first assessment classifying woodland by habitat type for each country and by region.

**Table 3.1** Area of woodland by habitat type for each country and region of GB

Region	Lowland beech/yew woodland	Lowland mixed deciduous woodland	Native pine woodlands	Non-HAP native pinewood	Upland birchwoods (Scot); birch dominated upland oakwoods (Eng, Wal)	Upland mixed ashwoods	Upland oakwoods	Wet woodland	Wood pasture & parkland	Broadleaf habitat NOT classified as priority	Non-native coniferous woodland	Clearfelled and transition	TOTAL
	Area (ha)	Area (ha)	Area (ha)	Area (ha)	Area (ha)	Area (ha)	Area (ha)	Area (ha)	Area (ha)	Area (ha)	Area (ha)	Area (ha)	Area (ha)
<b>GB</b>	<b>61,925</b>	<b>908,666</b>	<b>123,577</b>	<b>37,932</b>	<b>133,528</b>	<b>53,685</b>	<b>103,088</b>	<b>169,403</b>	<b>10,895</b>	<b>53,020</b>	<b>1,292,115</b>	<b>97,456</b>	<b>3,045,290</b>
<b>ENGLAND</b>	<b>54,482</b>	<b>747,508</b>	<b>0</b>	<b>0</b>	<b>11,220</b>	<b>31,907</b>	<b>44,027</b>	<b>77,789</b>	<b>7,730</b>	<b>19,401</b>	<b>327,536</b>	<b>21,967</b>	<b>1,343,568</b>
North West England	2,980	45,578	0	0	4,004	5,101	9,351	8,081	264	4,356	39,085	2,608	121,408
North East England	1,409	23,494	0	0	3,628	4,965	1,513	5,008	15	2,058	73,895	1,518	117,503
Yorkshire and the Humber	3,680	59,873	0	0	1,421	3,241	5,192	5,704	280	3,847	35,511	1,235	119,984
East Midlands	724	70,148	0	0	431	2,390	5,219	5,619	798	1,017	14,758	1,474	102,577
East England	3,739	94,807	0	0	0	309	840	13,497	1,178	1,101	34,368	4,468	154,305
South East England	24,700	245,410	0	0	0	1,758	323	9,761	4,946	2,740	49,670	5,582	344,890
South West England	15,133	138,219	0	0	1,336	8,337	15,586	23,212	154	2,459	54,579	3,021	262,036
West Midlands	2,116	69,980	0	0	401	5,805	6,004	6,906	95	1,824	25,671	2,060	120,863
<b>SCOTLAND</b>	<b>1,018</b>	<b>81,946</b>	<b>123,577</b>	<b>37,932</b>	<b>120,087</b>	<b>14,634</b>	<b>32,934</b>	<b>63,439</b>	<b>2,679</b>	<b>21,396</b>	<b>819,388</b>	<b>69,861</b>	<b>1,388,891</b>
North Scotland	0	4,034	31,642	16,626	25,848	1,269	4,764	9,243	0	2,007	106,962	21,110	223,503
North East Scotland	208	12,417	66,236	20,387	20,054	616	1,176	9,182	204	2,567	92,691	6,889	232,627
East Scotland	512	14,974	17,287	556	14,586	1,937	3,438	5,795	468	2,199	67,269	4,707	133,728
South Scotland	96	41,681	1,424	0	13,874	7,128	6,206	19,591	2,007	8,413	310,353	14,625	425,398
West Scotland	201	8,840	6,988	364	45,725	3,685	17,351	19,627	0	6,209	242,113	22,531	373,635
<b>WALES</b>	<b>6,424</b>	<b>79,211</b>	<b>0</b>	<b>0</b>	<b>2,221</b>	<b>7,144</b>	<b>26,126</b>	<b>28,175</b>	<b>486</b>	<b>12,224</b>	<b>145,192</b>	<b>5,628</b>	<b>312,831</b>

Note: 1. The wood pasture area in the above table only relates to wood pasture as defined by NFI as woodland; i.e. woodland must be over 0.5 hectares in extent, over 20% canopy cover and 20 m in width to qualify. Wood pasture can include land that contains less than 20% canopy cover. This estimate will therefore be a low estimate of the true value of wood pasture. 2. Area includes a revision to underlying total woodland area (this will be revised in Forestry Facts and Figures Sept 2020). Comparisons with woodland area should not be made as woodland area is defined differently to woodland habitat (with woodland area being greater than habitat area due to the treatment of land uses like open space and clear-fell). 3. Non-native woodland included for comparative purposes. 4. Non-native woodland may include PAWS sites.

# NFI woodland condition classifications

**Table 3.2** Area of woodland by woodland type for each country and region of GB

Region	Native	Near native & fragments	Non native	Not determinable	Total
	Area (ha)	Area (ha)	Area (ha)	Area (ha)	Area (ha)
<b>GB</b>	<b>1,507,105</b>	<b>56,776</b>	<b>1,461,267</b>	<b>19,629</b>	<b>3,044,777</b>
<b>ENGLAND</b>	<b>914,095</b>	<b>29,459</b>	<b>398,186</b>	<b>1,706</b>	<b>1,343,446</b>
North West England	73,932	2,329	45,152	421	<b>121,834</b>
North East England	36,201	3,005	77,982	149	<b>117,338</b>
Yorkshire and the Humber	68,954	3,783	46,837	271	<b>119,845</b>
East Midlands	73,964	2,305	26,272	37	<b>102,577</b>
East England	107,595	3,323	43,206	181	<b>154,306</b>
South East England	280,796	6,494	57,302	308	<b>344,901</b>
South West England	186,732	4,654	70,192	205	<b>261,782</b>
West Midlands	85,920	3,566	31,243	134	<b>120,863</b>
<b>SCOTLAND</b>	<b>442,611</b>	<b>20,313</b>	<b>908,259</b>	<b>17,205</b>	<b>1,388,388</b>
North Scotland	94,541	3,303	119,908	5,751	<b>223,503</b>
North East Scotland	111,260	3,781	115,197	2,248	<b>232,485</b>
East Scotland	48,860	2,647	80,049	2,171	<b>133,728</b>
South Scotland	78,739	7,026	336,526	2,709	<b>425,000</b>
West Scotland	109,211	3,556	256,579	4,327	<b>373,672</b>
<b>WALES</b>	<b>150,399</b>	<b>7,004</b>	<b>154,822</b>	<b>718</b>	<b>312,943</b>

Notes: 1. Woodland types are defined in Section 1.3.6. 2. Area includes a revision to underlying total woodland area (this will be revised in Forestry Facts and Figures Sept 2020). Comparisons with woodland area should not be made as woodland area is defined differently to woodland habitat (with woodland area being greater than habitat area due to the treatment of land uses like open space and clear-fell). Clear-fell woodland is included in the estimates, of which some cannot be classified and is set to not determinable.

This is the first assessment of the extent of native woodland and priority habitat type, across the whole of England, based on a balanced stratified random sample of all woodlands. There have been two previous assessments of native woodland extent; The Native Woodland Survey of Scotland (NWSS) and an analysis carried out using National Vegetation Classification (NVC) survey information. NWSS is a census of native woods, with a nominal date of 2012, which identified and mapped the location, extent, type and condition of all of Scotland's native and nearly native woods, as well as plantations on ancient woodland sites (PAWS). As two relatively recent assessments, the NFI and NWSS estimates broadly align, although some differences emerge due to the different methodologies used. In comparison, the 2008 JNCC created estimates of the area of priority habitat types used National Vegetation Classification (NVC) data (Rodwell 1991). Differences between these and the NFI results largely arise from the lower sample size and non-random sampling associated with the NVC dataset. In general, due to the methodological differences, direct comparisons between the three datasets should not be made.

## 4 Stand-level condition classification results for native woodland area in England

**Table 4.1** Area of native woodland stands in England classified as favourable, intermediate or unfavourable for each of the 15 WEC indicators

Native	Unfavourable		Intermediate		Favourable	
	Area (000 ha)	SE%	Area (000 ha)	SE%	Area (000 ha)	SE%
Ground flora	172.9	2.7	691.3	1.1	49.8	5.5
Tree pests and diseases	22.4	8.0	127.0	3.1	764.7	1.0
Invasive species	91.2	3.9	9.9	12.6	813.0	0.9
Herbivores / grazing pressure	345.6	1.8	113.0	3.1	455.5	1.6
Regeneration at component group level	0.0	–	737.6	1.0	176.5	2.6
Number of native tree and/or shrub species	101.2	3.8	192.9	2.5	620.1	1.1
Deadwood volume (m <sup>3</sup> per ha)	724.8	1.0	141.7	2.9	47.6	5.2
Vertical structure	61.9	5.0	312.3	2.0	540.0	1.2
Veteran trees	905.8	0.8	1.6	34.0	6.7	14.4
Age distribution of tree species	157.7	3.0	524.1	1.3	232.3	2.2
Nativeness of occupancy	0.0	–	129.3	2.9	784.8	0.9
Proportion of open space	763.8	0.9	137.7	3.0	0.9	28.2
Proportion of woodland / favourable habitat	122.1	3.5	346.5	1.9	445.5	1.6
Size of woodland parcel	224.0	2.8	209.0	2.8	481.1	1.4
Regeneration at population level	39.5	6.6	581.2	1.4	293.5	2.1
Overall ecological condition score	4.6	18.2	815.1	0.9	82.7	4.1


### Notes:

1. SE = standard error. Amber text = values with SE >25%.
2. Native is defined in Section 1.3.6.
3. Woodland ecological condition classification categories and indicators are defined in the methodology.
4. For more information refer to the methodology report.
5. Classification thresholds for ground flora; favourable, unfavourable and intermediate are adjusted for different woodland NVC types, with 'more' being expected of nutrient rich types than nutrient poor types.



# NFI woodland condition classifications

**Figure 4.1** The proportion of each woodland ecological condition class, for each WEC indicators in native woodland stands in England



**Notes:**

1. Native is defined in Section 1.3.6.
2. Woodland ecological condition classification categories and indicators are defined in the methodology.
3. For more information refer to the methodology report.
4. Classification thresholds for ground flora; favourable, unfavourable and intermediate are adjusted for different woodland NVC types, with 'more' being expected of nutrient rich types than nutrient poor types.

## 5 Stand-level condition classification results for near native woodland and fragments in England

**Table 5.1** Area of near native woodland and fragments in England classified by woodland ecological condition for each of the 15 WEC indicators


Near native & fragments	Unfavourable		Intermediate		Favourable	
	Area (000 ha)	SE%	Area (000 ha)	SE%	Area (000 ha)	SE%
Ground flora	3.9	17.3	23.7	6.7	1.8	24.7
Tree pests and diseases	0.6	43.6	3.8	18.0	25.1	6.5
Invasive species	3.5	17.2	0.2	81.2	25.7	6.5
Herbivores / grazing pressure	10.7	10.0	4.7	12.8	14.1	9.2
Regeneration at component group level	0.0	–	25.5	6.5	3.9	16.9
Number of native tree and/or shrub species	5.7	12.0	10.1	10.6	13.6	9.2
Deadwood volume (m <sup>3</sup> per ha)	23.3	6.8	4.1	16.0	2.0	24.2
Vertical structure	3.4	18.6	9.2	10.4	16.9	8.1
Veteran trees	29.2	6.1	0.0	–	0.3	71.0
Age distribution of tree species	6.5	13.0	19.6	7.4	3.4	17.5
Nativeness of occupancy	28.1	6.3	0.4	23.6	1.0	17.6
Proportion of open space	25.7	6.5	6.1	12.7	< 0.1	68.2
Proportion of woodland / favourable habitat	4.4	16.9	10.7	10.6	14.4	8.1
Size of woodland parcel	7.3	12.9	5.1	14.8	17.1	7.8
Regeneration at population level	1.8	29.8	19.1	7.7	8.6	10.3
Overall ecological condition score	1.3	30.3	30.3	5.9	0.2	82.3

### Notes:

1. SE = standard error. Amber text = values with SE >25%.
2. Woodland ecological condition classification categories and indicators are defined in the methodology report.
3. Near native and fragments are defined in Section 1.3.6.
4. For more information refer to the methodology report.
5. Classification thresholds for ground flora; favourable, unfavourable and intermediate are adjusted for different woodland NVC types, with 'more' being expected of nutrient rich types than nutrient poor types.
6. Size of woodland parcel: Fragments tend to occur in larger woods, therefore the results may reflect the overplanting of native woodland with large non-native plantation woods. NFI reports on woodland area (2012) evidence that non-native woods are larger on average than native woods.

# NFI woodland condition classifications

**Figure 5.1** The proportion of each woodland ecological condition class, for each WEC indicator type in near native woodland and fragments in England



**Notes:**

1. Woodland ecological condition classification categories and indicators are defined in the methodology report.
2. For more information refer to the methodology report.
3. Classification thresholds for ground flora; favourable, unfavourable and intermediate are adjusted for different woodland NVC types, with 'more' being expected of nutrient rich types than nutrient poor types.
4. NFI analysis has found that the fragments within near native tend to occur in larger non – native plantation woodlands.

## 6 Stand-level condition classification results for non-native woodland stands in England

**Table 6.1** Area of non-native woodland stands in England classified by woodland ecological condition classification for each of the 15 WEC indicators


Non native	Unfavourable		Intermediate		Favourable	
	Area (000 ha)	SE%	Area (000 ha)	SE%	Area (000 ha)	SE%
Ground flora	57.7	4.1	302.0	1.4	38.5	5.2
Tree pests and diseases	12.0	9.5	39.3	5.7	346.8	1.2
Invasive species	30.9	6.2	2.1	24.2	365.1	1.1
Herbivores / grazing pressure	117.8	2.8	65.0	3.6	215.4	2.0
Regeneration at component group level	0.0	–	358.8	1.2	39.4	5.3
Number of native tree and/or shrub species	234.4	1.6	93.1	3.3	70.7	4.0
Deadwood volume (m <sup>3</sup> per ha)	283.1	1.5	83.4	3.5	31.7	6.2
Vertical structure	100.9	3.1	183.2	2.1	114.1	3.1
Veteran trees	397.4	1.0	0.0	–	0.8	35.5
Age distribution of tree species	177.4	2.0	206.5	2.0	14.4	8.7
Nativeness of occupancy	398.2	1.0	0.0	–	0.0	–
Proportion of open space	336.6	1.3	70.8	3.7	0.7	29.2
Proportion of woodland / favourable habitat	30.1	6.6	94.2	3.5	273.8	1.4
Size of woodland parcel	49.2	5.2	45.7	5.3	303.3	1.2
Regeneration at population level	40.9	6.0	258.9	1.7	98.4	3.2
Overall ecological condition score	17.9	8.5	389.2	1.1	1.0	36.7

Notes:

1. SE = standard error. Amber text = values with SE >25%.
2. Non-native is defined in Section 1.3.6.
3. Woodland ecological condition classification categories and indicators are defined in the methodology report.
4. For more information refer to the methodology report.
5. Classification thresholds for ground flora; favourable, unfavourable and intermediate are adjusted for different woodland NVC types, with 'more' being expected of nutrient rich types than nutrient poor types.

# NFI woodland condition classifications

**Figure 6.1** The proportion of each woodland ecological condition class, for each WEC indicator type in non-native woodland stands in England




**Notes:**

1. 'Non-native' is defined in Section 1.3.6.
2. Woodland ecological condition classification categories and indicators are defined in the methodology report.
3. For more information refer to the methodology report.
4. Classification thresholds for ground flora; favourable, unfavourable and intermediate are adjusted for different woodland NVC types, with 'more' being expected of nutrient rich types than nutrient poor types.

## 7 Condition scoring distribution

**Figure 7.1** The overall distribution of ecological condition class by woodland type in England



Notes: 1. Native = native woodland area, Near native and fragments = Near native woodland area and fragments, non-native = non-native woodland area. 2. The NFI calculator is used to score each of the 15 ecological condition indicators that can then be combined and used to give an over-all score, and classification as favourable (fav) score 36-45, intermediate (int) score 26-35 or unfavourable (unfav) score 16-25 by woodland type. 3. Dashed line = threshold of each condition classification. To inform where to set the thresholds for each of the three classification categories published evidence was used. 4. Woodland types are defined in Section 1.3.6. 5. Refer to the methodology report for more information.

## 8 Discussion

This is probably the largest and most in-depth study of ecological condition of any habitat in England to date. All types of England's woodlands, native and non-native, semi natural and plantation have been measured in depth, to the same standard, using a statistically robust stratified random sampling method. The statistical results present a robust evidence base and source of reference data on which to build the scores. The scoring mechanism and classification thresholds have been developed over a number of years and represent a balanced and equal assessment of condition status.

Most of the scores presented are based on statistics derived from objective measures such as tree counts, deadwood diameters and species identification and as such are easily replicable and unambiguous. Other more subjective or difficult to identify measures, such as % levels of herbivore browsing damage, or % of squirrel stripping high in the canopy are harder to quantify and may under-report to some extent, but overall the broad trends identified hold and have stood up to testing against comparable data sources and expert opinion.

### 8.1 Woodland habitat area

One of the most notable results from this study was that the total area of native woodland in England is estimated to be 914 thousand hectares. This is circa 250 thousand hectares greater than the 660 thousand hectares previously estimated in the 2008 HAP assessment. This means that there is around 38% more woodland that can be classified as native in England than was previously reported. The majority of this newly identified native woodland occurs predominantly in lowland woods which are smaller than 2 ha in extent. The reason that the present study has identified these areas of native woodland, where previous studies have not, is because of advances in earth observation technology in recent years. Furthermore, the NFI, as compared to previous assessments, stratified out smaller stands of  $\geq 0.1$  hectares and  $< 0.5$  hectares, meaning simply that more small areas of native woodland habitat, situated within other woodland habitat types, have been identified and included in the study.

The increase in the estimate of native woodland area in England is reflected in estimates of the area of the various native woodland Habitat Action Plan (HAP) types. In this study, the NFI reduced the minimum area of a 'qualifying stand' of a woodland habitat within a woodland from  $\geq 0.1$  ha to  $\geq 0.01$  ha. Therefore, woodland types such as native wet woodland, which naturally occur in smaller localised patches (of wetter land), have been recorded and included, where previously they may have been 'lost' in the counts of other woodland types. Native wet woodland is now estimated to be 78 thousand hectares compared to circa 30 thousand hectares (recorded in the 2008 HAP estimates).

There has also been a large increase in the area classified as lowland mixed deciduous woodland. NFI now estimates 748 thousand hectares of lowland mixed deciduous

# NFI woodland condition classifications

---

woodland in England, which is higher than previously estimated in the 2008 HAP assessment. This increase is due, again, to advances in technology such as earth observation that mean it is now easier to identify smaller woodlands in the lowlands, where previously this was much more difficult.

## 8.2 Classification of woodland condition

Woodland condition was measured, then scored and classified, as either favourable, intermediate or unfavourable. The scoring method was to take each woodland stand in the survey and compare each of the woodland ecological indicators to the characteristics of a benchmark, namely 'semi-natural woodland in favourable condition', (refer to the *NFI woodland ecological condition in Great Britain: Methodology report* for more information). To inform where to set the thresholds for each of the three-classification categories, published evidence was used where available but in areas where published evidence was unavailable, expert opinion was used. For example, to determine a 'favourable' amount of seedlings, saplings and small trees in a woodland, a combination of published evidence and expert opinion was used to set the thresholds. Limiting the list of woodland ecological condition indicators to 15 measures is a simplification of an exceptionally complex ecosystem; however, it is a compromise between the feasibility of carrying out a study of this scale and achieving its objectives. Other similar studies that assess the ecological condition of habitats have used fewer than 15 measures; due to the absence of published data to inform thresholds (NWSS 2012: [forestry.gov.scot](http://forestry.gov.scot)).

Whilst the classification framework may be considered a simplified model of woodland ecology, it does provide a good indication of where potential issues may lie or where potential positive trends may occur. For example, volume of deadwood can be used as an indicator of biodiversity based on the assumption that the presence of deadwood provides an ecological niche used by fauna and flora. Therefore, a greater volume of deadwood is scored as favourable because it is assumed to increase biodiversity in woodlands. However, whilst this assumption holds true in the majority of instances there may be circumstances where it does not, such as in clear felling situations. Caution should therefore be applied to interpretation or inference drawn from these classification results, as they are indicative in nature as opposed to an absolute measure of ecological condition status. Whilst the model may not fit all individual situations or stands, analysis suggests that it can provide an overall statistically robust picture.

The application of a single "semi-natural" benchmark against all woodlands, which may be non-native plantations or coppice regimes, could be considered to be inappropriate because semi-natural status is not necessarily an objective of these woodland types. However, the application of a consistent 'semi-natural' benchmark to all woodland types (i.e. non-native plantations, coppice and native high forest) does provide a transparent and consistent measure allowing all woodland types to be compared equally.

This consistent application of a single benchmark has reflected positively on non-native plantations overall, which have scored reasonably well and more highly than may have been expected, with 96% scored as intermediate or favourable condition. In addition,



# NFI woodland condition classifications

---

only 4% of non-native woodland area was classified as unfavourable, and just under 0.5% of native woodlands classified as unfavourable. Plantations scored equally or better than some native woods for some of the ecological indicators, for example, plantations generally contain more deadwood than native woodlands. Furthermore, the higher scores are also likely to reflect how the plantations of the mid-20<sup>th</sup> century are maturing and diversifying with age, meaning the scores for the presence of natives, the number of stories and the diversification of age classes within non-native woodlands stand up positively to the benchmark.

These results provide useful insight into understanding the main physical and observable drivers behind woodland ecological condition in England today. The results suggest that the main issues are:

1. A lack of older trees and stands
2. Low numbers of veteran trees
3. Low volume of deadwood
4. Habitat fragmentation
5. Herbivore pressure and its impact on vegetation and regeneration
6. The notable presence of invasive and pest and diseases

The first four issues are evidenced by the high proportion of woodland area in unfavourable or intermediate condition for these factors. The last two do not have such high levels of woodland area in unfavourable or intermediate condition, but are considered of concern as they have high impact. Woodland management could make significant changes to improve condition scores, increasing open space, shrub layer and natural regeneration, more quickly than allowing natural processes alone to change the overall woodland structure.

The study has also highlighted some positive results for woodland ecological condition in England:

1. There is now evidence to classify more woodland in England as native than was previously estimated.
2. 99% of native woodlands are classified as either favourable or intermediate ecological status.
3. 86% of native woods achieved favourable status for native-ness of occupancy
4. 68% of native woods achieved favourable or intermediate status for number of native trees and shrubs species in the canopy.
5. Some habitats previously thought rare such as wet woodland are much more extensive than was previously estimated.

## NFI woodland condition classifications

---

6. The relatively high amount of woodland stands scoring as favourable or intermediate for tree regeneration, for both native and non-native woods.

### 8.3 Conclusion

The results present a valuable insight into the current ecological condition of woodland habitat in England and provides a baseline and reference data set for future monitoring work. Further information on the methodology and underlying statistics is available in the companion reports.

## 9 References

See Methodology Report.

## 10 Glossary

Word/phrase	Definition
Age class	A grouping of trees into specific age ranges for classification purposes. For the purposes of the "age distribution of trees" NFI WEC indicator, trees are grouped into three age classes: 0 – 20 years (Young); 21 - 150 years (Intermediate); >150 years (Old). For birch, cherry or Sorbus species: 0 - 20 years (Young); 21 - 60 years (Intermediate) >60 years (Old). Not applicable is used for stands without trees.
Ancient semi-natural woodland (ASNW)	Woodland which has been in continuous existence since 1600 (1750 in Scotland).
Area (forest/woodland)	Forest and woodland area can be defined in net or gross terms. Net area is the land actually covered by trees (in the National Forest Inventory that is to the drip line of the canopy). Gross area includes both the area covered by trees and the open spaces (<0.5 hectare) within (e.g. rides, glades, ponds).
Bark stripping	The removal of bark from trees by herbivores.
Biodiversity	Biodiversity represents 'all heritability-based variation at all levels of organisation, from the genes within a single local population, to the species composing all or part of a local community, and finally to the communities themselves that compose the living parts of the multifarious ecosystems of the world' (Wilson, 1997, p.1)
Broadleaves	Trees and shrubs that belong to the angiosperms (flowering plants) (as distinct from the gymnosperms that includes conifers). Most in the UK are deciduous and have laminar leaves (they do not have needles or cones) and a few, such as alder, have cone-like structures for their seeds which are not true cones. Sometimes referred to as 'hardwoods'.
Browsing	Herbivores feeding on tree buds, shoots and foliage.
Canopy	The mass of foliage and branches formed collectively by the crowns of trees.
Canopy cover	The percentage cover of the canopy across a defined area (e.g. NFI survey section or square).
Clear-felling	Cutting down of an area of woodland (if it is within a larger area of woodland it is typically a felling greater than 0.25 hectare). Sometimes a scatter or small clumps of trees may be left standing within the felled area.
Common Standards Monitoring (CSM)	The CSM approach was established during the 1990s by UK conservation agencies to describe the condition of protected sites, such as Sites of Special Scientific Interest (SSSI), in order to assess the effectiveness of conservation policies and practice.
Component (or sub-component)	Individual elements of the NFI survey component group. For example, each tree species will be recorded under a separate component, as will each habitat type if two habitats are intimately mixed (such as upland birchwood and wet woodland).
Component group	Homogeneous areas of the NFI survey that are too small (<0.05 ha) to practically map using Geographic Information System (GIS) software in the field, but with most of the same defining characteristics as a section. Component groups can be subdivided into components.

# NFI woodland condition classifications

Word/phrase	Definition
Condition	Shorthand for Woodland Ecological Condition.
Conifers	Trees and shrubs that belong to the gymnosperms, as distinct from the angiosperms that include broadleaves). Conifers mostly have needles or scale-like leaves and are usually evergreen. Sometimes referred to as 'softwoods'.
Convention on Biological Diversity (CBD)	A multilateral treaty to develop national strategies for the conservation and sustainable use of biological diversity.
Crown dieback	The death of branches within a tree's crown.
Deadwood	Non-living woody biomass not contained in the litter, either standing or lying on the ground (the NFI 'volume of deadwood' indicator does not include data on stumps).
Diameter at breast height (DBH)	The diameter on the stem of a tree at 'breast height', defined as 1.3 m from ground level.
Drip line	The drip line is the furthest tip of the widest branch in the crown; the last point from which the tree can drip if wet. If two treed sections have drip lines that cross over each other use the centre line of the cross over.
Earth observation	The collection of information about the physical, chemical, and biological systems of the planet via remote-sensing technologies.
Ecology	The relations of organisms to one another and to their physical surroundings.
Establishment	The formative period that ends once young trees are of sufficient size that, given adequate protection, they are likely to survive at the required stocking. This varies for species and according to environmental condition, but is typically from around five to twenty years.
EU Habitats Directive	The EU Habitats Directive (Directive 92/43/EEC) aims to promote the maintenance of biodiversity by requiring Member States to take measures to maintain or restore natural habitats and wild species listed on its Annexes to a favourable conservation status (JNCC, 2018).
(Vegetation) field layer	Vegetation 10 cm to 2 m tall measured as part of the NFI vegetation assessment.
Flora	The plants of a particular region, habitat, or geological period.
Forest (or woodland)	See woodland
Forestry Commission (FC)	The government department responsible for regulating forestry, implementing forestry policy and managing state forests in England. It was formerly also responsible for Forestry in Wales and Scotland, however on 1 April 2013 the Forestry Commission's functions in Wales transferred to a new organisation, Natural Resources Wales. From 1 April 2019, forestry was fully devolved, except for common issues addressed on a GB or UK basis, such as international forestry, plant health and forestry standards. Following devolution, two new Scottish Government agencies were created, Scottish Forestry and Forest & Land Scotland.
Forestry and Land Scotland (FLS)	The Scottish Government agency responsible for managing Scotland's national forests and land.
Fragments	Small areas of woodland with 50% or more native tree species occupancy in the upper canopy, but that fall in the size range 0.05 ha to 0.099 ha.
Geographic Information System (GIS)	A system designed to capture, store, manipulate, analyse, manage, and present spatial or geographic data.

# NFI woodland condition classifications

<b>Word/phrase</b>	<b>Definition</b>
Global Positioning System (GPS)	A satellite-based global navigation satellite system that provides geolocation and time information to a GPS receiver.
Great Britain (GB)	England, Scotland and Wales.
(Vegetation) ground layer	Vegetation 0 – 10 cm tall measured as part of the NFI vegetation assessment.
Habitat Action Plan (HAP)	For all UK BAP priority habitats classified between 1995 and 1999, a Habitat Action Plan (HAP) was created (45 in total). For the habitats added to the priority habitats list in 2007, no UK action plans have been, or will be, produced, as conservation action is now primarily carried out at a country-level, rather than a UK-level, in response to the generation of country-level biodiversity strategies and aims (JNCC, 2019a).
Herbivore	An animal that is adapted to eating plant material for the main component of its diet.
Hectare (ha)	Unit of area defined as 10,000 square metres (100 m by 100 m), approximately equivalent to 2.47 acres.
Indicator	A quantitative or qualitative parameter that synthesises complex information and can be periodically measured to assess trends over time. 15 stand level indicators were selected to assess the condition of woodlands as part of the NFI WEC approach.
Invasive species	A species that is not native to a location, where it is likely to cause ecological or economic harm.
Invertebrate	A cold-blooded animal that does not have a backbone.
Lichen	A composite organism that arises from algae or cyanobacteria living among filaments of multiple fungi species in a mutualistic relationship.
National Forest Inventory (NFI)	National forest inventories are carried out in GB by the FC to provide accurate, up-to-date information about the size, distribution, composition and condition of the forests and woodlands. The current NFI, which began in 2009, is a multi-purpose operation that has involved the production of a forest and woodland map for GB and a continuing programme of field surveys of the mapped forest and woodland areas.
National Forest Inventory map	An earth observation-based programme that monitors and maps the extent and location of woodlands across GB on an annual basis.
National Forest Inventory field survey	A field survey of a large, stratified-random sample (15,100 sites) of woodlands across GB on a 5-year rolling cycle using a standardised protocol.
Native species	Species that have arrived and inhabited an area naturally, without deliberate assistance by man. For trees and shrubs in the United Kingdom usually taken to mean those present after post-glacial re-colonisation (around 11,000 years ago) and before historic times. Some species are only native in particular regions - hence locally native.
Natural England (NE)	The government's adviser for the natural environment in England. Natural England is an executive non-departmental public body, sponsored by Defra.
Naturalised species	A species that, once it is introduced outside its native distributional range, establishes self-sustaining populations.
Natural Resources Wales (NRW)	The organisation responsible for advising the Welsh Government on the environment, created on 1 April 2013. NRW is responsible for the functions previously carried out by the Environment Agency in Wales, the Countryside Council for Wales and Forestry Commission Wales.
National Vegetation Classification (NVC)	Vegetation classification system commonly used in Great Britain.

# NFI woodland condition classifications

Word/phrase	Definition
Native woodland	Woodland with 50% or more native tree species occupancy in the upper canopy that either: <ul style="list-style-type: none"> <li>- Forms a discrete woodland parcel with a minimum area of 0.5 ha.</li> <li>- Forms a woodland stand with a minimum area of 0.1 ha that is part of a woodland that is 0.5 ha or larger.</li> </ul>
Native Woodland Survey of Scotland (NWSS)	A survey of all native woodlands, nearly native woodlands and non-native plantations on ancient woodland sites in Scotland.
Near native woodland	'Nearly' native woodland with 40% to 49% native species canopy cover.
NFI Condition Calculator	An analytical GIS tool developed to automatically produces the component group-level NFI WEC indicator results per woodland type and aggregated statistics for the reporting area.
NFI WEC working group	The expert committee that was established to develop the NFI WEC indicator approach. This group consists of representatives from (former) FC England and Scotland, Scottish Natural Heritage, Natural England, Natural Resources Wales and the Welsh Government.
Non-native woodland	Woodland with less than 40% native species occupancy.
(Woodland) parcel	Discrete blocks of woodland that are separated from other woodland parcels by gaps of at least 20 m in length.
Private sector estate	Forests and woodlands in the UK not managed by the Forestry Commission, Natural Resources Wales or Forest Service. In the context of the National Forest Inventory, 'Private sector' is used for convenience although it includes land owned or managed by bodies such as local authorities and charities.
(Natural) Regeneration	The regeneration of existing woodland by natural means, i.e. without sowing or planting.
(Ecological) resilience	The ecological resilience of woodland ecosystems refers to their ability to absorb disturbance while maintaining the major habitat-forming species that define their structure and ecosystem functioning. Resilience incorporates both the woodland ecosystem's ability to resist changes in response to disturbance or, failing this, its capacity to recover functioning via adaptation.
(NFI) sample square	The one-hectare (100 m by 100 m) square plots, which may be entirely within woodland or may overlap the woodland edge, used for the NFI field survey.
Sapling	Young tree $\geq 50$ cm tall and $< 4$ cm in diameter.
Saproxyllic	Dependent on deadwood.
(NFI WEC) score	An ordinal score is assigned to the individual indicator classes of 'unfavourable' (1), 'intermediate' (2) and 'favourable' (3). The scores are summed for all 15 indicators to provide each stand's overall ecological condition score, which has a maximum value of 45.
Scottish Natural Heritage (SNH)	The public body responsible for protecting and promoting Scotland's natural heritage, especially its natural, genetic and scenic diversity. To be renamed 'NatureScot' from May 2020.
Section	Within each NFI sample square, the forest was stratified into different woodland 'sections'. Sections are defined by individual strata at least 0.05 ha in size that are differentiated on basis of forest type, habitat, land use, silviculture system, tree and shrub composition, age and structure.
Seedling	Young tree $< 50$ cm tall.
Shrub	Woody plants often (but not always) branching abundantly from the base that are between 2-5 m tall.

# NFI woodland condition classifications

<b>Word/phrase</b>	<b>Definition</b>
(Vegetation) shrub layer	Vegetation 2-5 m tall measured as part of the NFI vegetation assessment.
Site of Special Scientific Interest (SSSI)	A formal conservation designation that is applied to areas of particular interest to science because of the geology/geomorphology features or species it contains or supports.
(Woodland) Stand	A distinct area of woodland (from either planting or natural regeneration), generally composed of a uniform group of trees in terms of species composition and spatial distribution, and age and size class distribution.
Standard error (SE)	The measure of the margin of error associated with an estimate as a result of sampling from a population with statistical variability. Larger standard errors indicate less precision in the estimate. Standard errors in this report are quoted in relative terms (i.e. as percentages of the value of the estimate). In this report, any standard error greater than 25% is reported in amber italics and represents a lower degree of assurance in the estimates.
Stocking	The density of trees within a woodland.
Stump	The above-ground base part of a tree that would usually remain after felling.
Transect	A path along which a survey is carried out.
Transition woodland	Land classified as woodland area that is in transition between no tree cover and tree cover. Examples include clear-fell sites, restock sites, new planting sites and land with natural regeneration.
UK Biodiversity Action Plan (UK BAP)	The UK government's national biodiversity action plan that was developed in response to the Convention on Biological Diversity and replaced by the 'UK Post-2010 Biodiversity Framework' in 2012 following new international targets. It described the biological resources of the UK and provided detailed plans for conservation of these resources (JNCC, 2019a). The UK BAP priority habitats were identified as the most threatened habitats requiring conservation action under the UK BAP.
UK Forestry Standard (UKFS)	The reference standard for sustainable forest management across the UK that applies to all woodland to ensure that international agreements and conventions on areas such as sustainable forest management, climate change, biodiversity and the protection of water resources are applied in the UK.
United Kingdom (UK)	Great Britain and Northern Ireland.
Woodland (or forest)	Land predominately covered in trees (defined as land under stands of trees with a canopy cover of at least 20%, or the ability to achieve this, and with a minimum area of 0.5 hectare and minimum width of 20 m), whether in large tracts (generally called forests) or smaller areas known by a variety of terms (including woods, copses, spinneys or shelterbelts).
(NFI) Woodland Ecological Condition (WEC)	The approach used by the NFI to assess the ecological condition of woodlands in GB in terms of their likely biodiversity value.
(Woodland) storey	A woodland's trees and shrubs can often be stratified into distinct layers, or storeys, according to their height.
Vertical (woodland) structure	The number of canopy storeys present.
Veteran trees	A tree that is of interest biologically, culturally or aesthetically because of its age, size or condition (Read, 2000)



## 11 Annex

**Table 11.1** Classification threshold summary

Woodland Ecological Condition Indicator	Classification thresholds (abbreviated)		
	Favourable condition	Intermediate condition	Unfavourable condition
Age distribution of trees	Three age classes present	Two age classes present	One age class present
Wild, domestic and feral herbivore damage	No herbivore or squirrel damage	Herbivore damage near to stand Squirrel damage only	Herbivore damage in stand
Invasive plant species	No invasive species in stand	Rhododendron and laurel not present, other invasive species < 10% cover	Rhododendron and laurel present, or other invasive species > 10% cover
Number of native tree species	Five or more native tree or shrub species in a stand	Three to four native tree or shrub species in a stand	None to two native tree or shrub species in a stand
Occupancy of native trees (in a stand)			
England and Wales	> 80%	50-80%	< 50%
Scotland	>90%	80-90%	<50%
Open space within woodland			
Woods greater than 10 hectares	10-25% open & ≥ 50% of open high quality	<10% open space, yet over 50% of open high quality >25 and <50% open space & > 50% high quality open space	≥ 50% open space, irrespective of quality of open <10% open space and over 50% of open space low quality
		10-25% open space and over 50% of open space low quality	
Woods less than 10 hectares	0-10% & > 50% of open high quality	10-25% open space with over 50% high quality open 0-10% open space and less than 50% of open space high quality	>25% open space 10-25% open space and over 50% low quality open
Proportion of favourable land cover around woodland	>20%	10-20%	<10%
Woodland Regeneration: Stand	All three classes present in stand; Trees 4-7cm dbh, saplings and seedlings	One or two classes only present in stand	Not applicable
Woodland Regeneration: Square-level	All three classes present nearby; Trees 4-7cm dbh, saplings and seedlings	One or two classes only present nearby	No classes present nearby
Tree health	Tree mortality less than 10%, no pests or diseases and no crown dieback	11% to 25% mortality	Greater than 25% tree mortality
		0-11% mortality and crown dieback or low risk pest or disease present	Any high risk pest or disease
Vegetation and ground flora	Strong plant community	Average plant community	Weak plant community
Woodland vertical structure	Four or more storeys in a stand or a complex stand	2 or 3 storeys in a stand	One or less storey in a stand
Veteran trees	Greater than 40 veteran trees per 20 hectares	Between one and 40 veteran trees per 20 hectares	Less than 1 veteran tree per 20 hectares
Volume of deadwood	Greater than 80 cubic metres per hectare	Between 20 and 80 cubic metres per hectare	Less than 20 cubic metres per hectare
Size of woodlands	Greater than 20 hectares	between 5 and 20 hectares	less than 5 hectares
Score per WEC indicator	3	2	1
Total score	Greater than 35	26 to 35	Less than 25





## 12 NFI national reports

This inventory report is one of a series of publications reporting the outputs of the Forestry Commission National Forest Inventory.

These and NFI data can be found on the NFI website [www.forestresearch.gov.uk/inventory](http://www.forestresearch.gov.uk/inventory).

### Official Statistics

This is not an Official Statistics publication, but it does include two tables of official statistics (Tables 3.1 and 3.2). More information about Official Statistics and the UK Statistics Authority is available at [www.statisticsauthority.gov.uk](http://www.statisticsauthority.gov.uk)

National Forest Inventory Statistician: David Ross

### Survey design

B. Ditchburn and A. Brewer

### Authors

B. Ditchburn, T. Wilson, L. Henderson, K. Kirby and P. Steel.

### Publication support

V. Correia, D. Ross, W. Peden, H. Stephenson, L. Halsall and C. Bellamy

### Reviewers and steer

For a list of those who assisted in the quality assurance of these statistics, please see the Forest Research [statistics webpages](#).

Lead Reviewers: Colin Edwards, Emma Goldberg, Jeanette Hall, Rebecca Isted, Fiona Macfarlane, Neil Riddle and Chris Tucker

For more information and to view and download Forest Research publications, visit: [www.forestresearch.gov.uk/publications](http://www.forestresearch.gov.uk/publications)

We will consider all requests to make the content of our publications available in alternative formats. Please email: [publications@forestresearch.gov.uk](mailto:publications@forestresearch.gov.uk)