

# The Integrated Management of *Hylobius abietis* in UK Forestry



Dr Ian H. Willoughby, Dr Roger Moore and Dr Tom R. Nisbet

The Research Agency of the Forestry Commission



Forest Research is the Research Agency of the Forestry Commission and is the leading UK organisation engaged in forestry and tree related research. The Agency aims to support and enhance forestry and its role in sustainable development by providing innovative, high quality scientific research, technical support and consultancy services.



Figure 1. Adult Hylobius abietis browsing on a young conifer tree



## Contents

Summary1
1. Aims2
2. Background2
3. Management options3
3.1. Integrated forest management 3   3.2. Take no action 4   3.3. Avoid the problem 4   3.3. Avoid the problem 4   3.3.1. Alternative silvicultural systems or species 4   3.3.2. Fallow ground strategy 7   3.3.3. Manipulating ground vegetation 9   3.4. Using good quality planting stock 9   3.4. Take remedial action 9   3.4.1. Biological control with entomopathogenic nematodes 9   3.4.2. Mulching and de-stumping 10   3.4.3. Physical protection 11   3.4.4.1 Pre-planting pre-treatment with alpha-cypermethrin (Alpha C 6ED®) 12   3.4.4.1 Pre-planting pre-treatment with acetamiprid (Gazelle SG®) applied via 13   3.4.2. Pre-planting pre-treatment with acetamiprid (Gazelle SG®) applied via 14   3.4.4.3 Immediate post-planting spraying with cypermethrin or acetamiprid
4. Conclusions23
5. References
6. Appendix - Acetamiprid safety29
6.1. Introduction296.2. The approvals process for acetamiprid306.3. Some hazard and risk characteristics of acetamiprid31

### List of Tables

Table 1. Typical peak <i>Hylobius</i> damage periods, and possible safe planting windows,	
when following a fallow ground strategy	.8

Table 2. A summary of the relative costs, efficacy and potential risks of the mainremedial control measures available for protecting trees from damage by *Hylobius*abietis18

Table 3. A comparison of some of the hazard characteristics of insecticides used forprotecting trees from damage by *Hylobius abietis*22



## Summary

The large pine weevil (*Hylobius abietis* L.) is probably the most serious pest of newly planted or naturally regenerating woodland trees on restocking sites in the UK. On affected sites, in the absence of protective measures, losses of replanted trees will average around 50%, but in the worst cases all trees can be killed. Both conifers and broadleaves are attacked and in some circumstances it can become impossible to establish trees and sustainably regenerate forests.

It is recommended that an integrated approach to the management of this pest is adopted. As described in this Report, this involves firstly understanding the life cycle and likely impacts of *Hylobius*, and then considering the full range of potential approaches available to prevent the insect damaging young trees, and finally selecting a restocking management strategy that may involve single or multiple techniques in combination with each other.

Although research into non-chemical approaches is ongoing, currently on many sites in the UK and Ireland it is likely that insecticides will still need to be used as part of the integrated management of *Hylobius*. Where insecticide use is unavoidable, pre-treatment of young trees in an off-site horticultural tree nursery with alpha-cypermethrin or acetamiprid, possibly combined with later post-planting top-up sprays in the forest of those trees with acetamiprid, when used as described on the Health and Safety Executive approved pesticide product labels, and when combined with the additional precautionary measures in place as described in this Report, should not pose any unacceptable risk to consumers, operators, bystanders, neighbours or the wider environment.



## 1. Aims

The purpose of this Report is to provide updated guidance on the integrated pest management of the insect *Hylobius abietis*. Guidance on the integrated management of all major pest, weed and disease issues facing UK forestry, along with advice on pesticide minimisation and reducing the risk of damage to the environment, is provided in Forestry Commission Practice Guide 15: *Reducing pesticide use in forestry* (Willoughby *et al.*, 2004). This Report provides a summary of the main changes affecting integrated management of *Hylobius*. In doing so it draws on some of the key findings of an ongoing programme of collaborative forest industry research into alternatives to the use of the conventional insecticides for protecting trees from *Hylobius* damage (Hardy *et al.*, 2020; Willoughby *et al.*, 2020; Moore *et al.*, 2021). In addition, as a separate Appendix, it summarises knowledge and guidance on the safe use of the insecticide acetamiprid, which is increasingly being phased in across the UK forest industry as an alternative to cypermethrin.

## 2. Background

The large pine weevil (*Hylobius abietis* L., hereafter referred to as '*Hylobius*') is probably the most serious pest of newly planted or naturally regenerating woodland trees on restocking sites in the UK (Figure 1). Adult *Hylobius* are large insects - up to 12 mm long - and can live for up to 4 years in upland areas, or can complete their life cycle in as little as a year on lowland pine sites. The adult weevils migrate onto felled sites, either in low numbers if the felling is entirely new within a given forest area, or in higher numbers if the site is in close proximity to an existing clearfell. Eggs are laid below ground level in the bark of stumps of recently felled conifers or in fallen / felled conifer stems and branches in direct contact with the ground. Depending on the tree species, stumps can remain suitable for weevil reproduction for up to 4 years after felling. Climate also plays its part and in upland areas larval development in the stumps can take up to 2-4 years, but on lowland pine sites they can complete this development in as little as a year.

Larvae feed in the cambial layer inside the outer bark and pass through several moults before finally pupating in a pupal chamber (Figure 2). Adults usually emerge in the autumn and concentrate their feeding on the bark of exposed lower stems of transplanted or naturally regenerated conifer seedlings and the upper part of the stem of broadleaved trees. This often leads to girdling of the stem and results in plant death. Feeding is non-specific in the sense that adult weevils are known to browse on many available plant food sources including the branches of mature standing trees, seedlings, and woody weeds on site.

Adult *Hylobius* feed at any time of year when it is warm enough for insect activity, generally between early March and November. However, there is a tendency for two peaks of damage to occur, one in spring before egg laying and the other in late summer



soon after a new generation emerges from the stumps but before the adults hibernate underground. The timing of these feeding periods depends on both the physiological condition of the insects and the ambient temperature. The relative magnitude of these peaks will vary from forest to forest and from year to year. When sites are clearfelled the availability of alternative food sources for adult *Hylobius* is restricted, hence newly planted or naturally regenerating trees tend to be favoured by the weevils, giving rise to extensive browsing.

On affected sites, in the absence of protective measures, losses of replanted trees will average around 50%, but in the worst cases all trees can be killed. Both conifers and broadleaves are attacked and in some circumstances it can become impossible to establish trees and sustainably regenerate forests (Willoughby *et al.*, 2004).

## 3. Management options

### 3.1. Integrated forest management

Pesticide use in forestry is on a very small scale compared to agricultural, horticultural and amenity use. It has been estimated that despite forests covering over 13% of the land area of Britain, they account for less than 0.1% of all pesticides used (Willoughby *et al.*, 2009; McCarthy *et al.*, 2011).

Rather than using pesticides as a first resort, Forest Research recommend that an integrated approach to pest, disease and weed management (often referred to as 'IPM') is always adopted, based on an understanding of the nature and impacts of the damaging agent, and after consideration of the full range of potential solutions, if necessary in combination with each other. Only if all non-chemical control options have been considered and shown to be impractical, ineffective, excessively costly, or likely to carry the risk of causing more harm to people and the environment, should the use of pesticides be countenanced. Often the combination of non-chemical and chemical approaches can help to reduce the overall level of pesticide use on a given site. The core decision key (Figure 3) is taken from Forestry Commission Practice Guide 15 (Willoughby et al., 2004), and provides a simple means of adopting the integrated approach as recommended by Forest Research, thereby minimising pesticide use as required by the UK Forest Standard (Forestry Commission, 2017), and fulfilling the requirements of voluntary independent certification schemes such as those of the Forest Stewardship Council (FSC, 2019) or PEFC (PEFC, 2020), achieved through the UK Woodland Assurance Standard (UKWAS, 2018). The management options outlined in this Report are therefore grouped under the standard headings referred to in the core decision key.

The *Hylobius* Management Support System (Forest Research, 2021) is a decision support tool developed by Forest Research that utilises 'billet counts' to measure insect presence in combination with a model based on an understanding of *Hylobius* population dynamics to predict if damage is likely on a site, and hence what action or treatment is required. It



can be used to determine the optimum use of many of the approaches outlined in this Report under the headings of 'Avoid the Problem' and 'Remedial Control Measures', in particular the fallow ground strategy, choice of most suitable type of transplants, and the effects of entomopathogenic nematodes. Rather than simply relying on monitoring tree damage and when it occurs attempting to react to it quickly enough to prevent tree death, the system can help predict in advance where and when remedial action, in the form of post-planting spraying with insecticide (referred to as 'top-up spraying'), may be necessary. In effect the *Hylobius* Management Support System provides a sophisticated and effective means of following the core decision key, taking an integrated approach to the management of *Hylobius*, and minimising unnecessary pesticide use.

For lowland pine sites where *Hylobius* only has a one-year life cycle, a possible alternative approach to using the *Hylobius* Management Support System is to monitor weevil development in stumps through 'stump hacking' to predict the time of adult emergence and approximate population size. Depending on the risk and likely intensity of attack, sites can be left fallow for a period before planting, immediately restocked with treated trees, or immediately restocked with untreated trees. Monitoring of feeding damage on planted trees can then be used to identify recently planted areas that require top-up spraying. Full details and practical guidance on the use of this approach for lowland pine sites is given in Wainhouse *et al.* (2007). However, the use of stump hacking to monitor *Hylobius* populations in this way is likely to be considerably more expensive than the use of billet counts if a similar level of accuracy is required.

### 3.2. Take no action

With no action, losses of replanted trees will range from 0-100%, depending on *Hylobius* population levels, plant characteristics such as root collar diameter, availability of alternative food sources, and environmental conditions. Without monitoring of populations to allow decisions on intervention to be taken using techniques such as the *Hylobius* Management Support System, or very frequent monitoring of damage levels, the 'take no action' option is a very high-risk strategy and is therefore not recommended.

## 3.3. Avoid the problem

#### 3.3.1. Alternative silvicultural systems or species

For existing conifer forests, using alternative silvicultural systems to clearfelling and replanting with conifers can partly or completely avoid the problem of *Hylobius* damage to young trees. Conifer forests managed on a continuous cover basis are likely to have smaller areas of felling and more alternative food sources for emerging weevils, which means that naturally regenerating or planted trees can sometimes be established without the need for any further protection (Mason *et al.*, 2004). However, successful conversion to continuous cover forestry can take many years, and given the prevailing



windy, oceanic climate and inherent lack of tree stability on many British sites, such silvicultural systems are only ever likely to be appropriate for a minority (< 25%) of upland conifer plantations even in the long term (Paterson, 1990).

Although broadleaves planted on ex-conifer sites are liable to be attacked, once they are established, large or isolated broadleaved forests are unlikely to be vulnerable to *Hylobius* damage when felled areas are being restocked.

#### Figure 2. Hylobius abietis larva





#### Figure 3. The Core decision key (adapted from FC Practice Guide 15; www.forestresearch.gov.uk)





#### 3.3.2. Fallow ground strategy

Depending on the tree species, stumps can remain suitable for weevil reproduction for up to 5 years after felling. During this time two to three generations of *Hylobius* will emerge on site and, in the absence of suitable further conifer stumps to breed in, will tend to migrate to other sites. Allowing a clearfell site to lie fallow for at least 5 years may therefore allow trees to be planted with little risk of damage. A shorter period of fallow may be possible in some circumstances. See Table 1, (derived from Moore, 2004) for further details.

However, this approach will only work if neighbouring areas cannot act as sources of weevils. Hence freshly felled sites, or sites felled within the last 4 or 5 years, must be far enough away that the *Hylobius* that emerge from them, or are attracted to them, cannot travel to the newly planted ex-fallow site, but these requirements are not always practical. The safe separation distance between freshly felled and newly planted ex-fallow sites is around 1,000 m. For sites felled within the last 4 or 5 years, the safe separation distance is around 25-50 m. In this scenario planting a 25–50 m barrier strip of insecticide treated trees along the edge of the previously fallow site that is closest to the neighbouring felled area, should be sufficient to prevent significant damage to the untreated trees on the reminder of the planting site.

Adopting a fallow ground strategy can have adverse consequences on some sites. Long fallow periods can increase the risk of nitrate release, with potential impacts on stream acidity within acid sensitive and nitrogen saturated areas of the UK. Phosphate leaching may also increase leading to nutrient enrichment of drainage waters. Another possible impact on water is the extended loss of the water use benefit of forests for reducing downstream flood risk. On more fertile sites, leaving areas fallow allows competing vegetation to flourish, and this can dramatically increase the amount of herbicide which may need to be used to allow trees to establish, as well as necessitating the use of more intensive and disruptive cultivation techniques. Therefore on more fertile sites, adopting a fallow period strategy is likely to result in an increase in overall levels of pesticide use and significantly increased costs. Maintaining open areas within a forest block may also increase the risk of windthrow in adjacent stands. A wider consideration is that delaying planting for several years reduces carbon sequestration, and can also result in significant financial loss due to the revenue forgone, which is often not acceptable to the landowners.

Leaving sites fallow for at least 5 years is therefore only worth considering on less fertile, relatively sheltered clearfell sites that are not in sensitive riparian catchments, provided that the sites are far enough away from the nearest 0- to 5-year-old felling. Even then, leaving areas fallow is a high-risk strategy unless the site has been assessed using specific monitoring techniques, such as via the *Hylobius* Management Support System, and has been found or is predicted to have a sufficiently low *Hylobius* population.



#### Table 1. Typical peak Hylobius damage periods, and possible safe planting windows, when following a fallow ground strategy

Date of felling <sup>1</sup>	Initial damage period ( <i>Hylobius</i> site colonisation and egg laying) <sup>2,3</sup>	Peak damage periods ( <i>Hylobius</i> stump emergences) <sup>3</sup>	First safe planting date for treated trees, no top-up spraying <sup>4</sup>	First safe planting date for untreated trees, no top-up spraying <sup>5</sup>
Year 0, January – March	Year 0, April – August	Year 1, August – October	Year 4, spring	Year 5, spring.
		Year 2, April – June; August – October		(or back-end Year 4)
		Year 3, April – June		
Year 0, April – mid August	Year 0, April – August	Year 1, August – October	Year 5, spring	Year 6, spring
	Year 1, April – August	Year 2, April – June; August – October		(or back-end Year 5)
		Year 3, April – June; August – October		
		Year 4, April – June		
Year 0, mid-August – December	Year 1, April – August	Year 2, August – October	Year 5, spring	Year 6, spring
		Year 3, April – June; August – October		(or back-end Year 5)
		Year 4, April – June		

#### Notes

This table, which is derived from Moore (2004), gives an approximate indication of likely damage periods by *Hylobius*, and hence the periods when it may be safe to plant trees without the need to make post-planting (top-up) insecticide sprays if pursuing a fallow ground strategy. For a reliable assessment of the risk of damage, on-site *Hylobius* population levels and attributes of nearby felling sites (such as the area, felling dates, climate, species, pattern and location) must be taken into consideration. This can be done by the use of the *Hylobius* Management Support System (Forest Research, 2021).

- <sup>1</sup> Year 0 is taken to be the calendar year in which felling of the preceding crop takes place.
- <sup>2</sup> Damage to trees can be just as severe during this initial period of colonisation as the 'peak damage period'.
- <sup>3</sup> The period when damage will generally occur if the site is planted as soon as possible after felling.
- <sup>4</sup> The earliest date at which insecticide treated trees can be generally safely planted with no subsequent need for top-up insecticide sprays.
- <sup>5</sup> The earliest date at which untreated trees can generally be safely planted with no subsequent need for top-up insecticide sprays.



### 3.3.3. Manipulating ground vegetation

In the UK and Ireland, adult *Hylobius* move within clearfell sites mainly by walking, and there is some indication that they are discouraged from stopping to feed on trees in areas of open, weed-free ground with exposed mineral soil. In addition, areas with quite high weed cover of other woody plants, such as bramble (*Rubus fruticosus* agg. L.), can act as alternative food sources, thus diverting attacks away from tree seedlings. Hence the retention of a small amount of cover from non-invasive woody weeds, while maintaining open areas of mineral soil immediately surrounding tree seedlings by appropriate initial site cultivation followed by good weed control, which are in any case both standard elements of good silvicultural practice for tree establishment, may also help to reduce *Hylobius* browsing levels.

However, if large populations of weevils emerge they can overwhelm such defensive strategies and result in high losses of planted or naturally regenerated seedlings. In addition, leaving invasive weeds such as bramble unmanaged on site can itself kill young trees. Therefore, whilst this technique may make a useful contribution to reducing overall damage levels as part of an integrated approach to management, it should not be relied on in isolation.

### 3.3.4. Using good quality planting stock

Seedlings that are vigorous with good root systems and high quantities of resin are better able to withstand attack than small plants with poor defensive systems. All planting stock used should therefore have a good balance between roots and shoots and should conform to British Standard 3936 (Morgan, 1999), as this is also good silvicultural practice and helps to reduce the establishment period and the amount of subsequent weeding inputs required. The larger the diameter, the thicker the bark and the more resin that a tree seedling produces, and the more likely it is to be able to tolerate low to moderate levels of browsing by *Hylobius*. In Scandinavian work, trees with root collar diameters of 10mm or more have been shown to tolerate insect browsing damage without dying (Thorsen *et al.*, 2001). However, initial attacks can attract other adult weevils to the damaged plants and, if pest populations are high as they often are in the UK, the plant will quickly succumb to multiple attacks. For this reason, whilst the use of good quality, well balanced planting stock with larger root collar diameters may make a useful contribution to reducing the overall impact of *Hylobius*, as part of an integrated approach to management, it should not be relied on in isolation.

## 3.4. Take remedial action

A summary of the main remedial control options is given in Table 2.

#### 3.4.1. Biological control with entomopathogenic nematodes

Research has shown that the application of entomopathogenic nematodes (*Steinernema carpocapsae* Weiser) to late larval and pupal stages of *Hylobius* in stumps can reduce the



numbers of emerging adults, resulting in overall population reductions of up to 20% on spruce sites and between 70-90% on pine sites (Brixey *et al.*, 2006). This leads to fewer adults moving on to nearby restocking sites, and if nematodes are applied to consecutive clearfells within the same forest block, the total population pressure from *Hylobius* can be reduced to non-damaging levels. When combined with predictions of post-felling population dynamics derived from the *Hylobius* Management Support System, applications of nematodes can therefore help to achieve a significant reduction in the need to use insecticides.

High volumes of water must be used to apply the nematodes to tree stumps, and specialist forwarder mounted spray rigs are required. This makes nematode use impractical in remote locations where transport of water is difficult and expensive, or in situations where forwarder machinery cannot be used (such as on steep sites, or on soft ground if the brash mats used to protect the soil from wheeled or tracked machinery during harvesting operations have degraded). These sorts of conditions are relatively common in upland conifer plantations. In addition, nematodes are less effective where adult *Hylobius* can migrate from neighbouring forests under different ownership and management, or in forest blocks where clearfelling is small scale, sporadic or isolated, which can often be the case in many smaller, lowland forests.

In summary whilst biological control with entomopathogenic nematodes may have a valuable role to play as part of an integrated approach for managing the problem of *Hylobius* damage in some situations, it is estimated that between a third and two thirds of restocking sites in the UK will never be suitable for the technique due to the limitations of site geography, even if the application technology can be improved and costs can be reduced in the future.

#### 3.4.2. Mulching and de-stumping

Mulching or de-stumping have the potential to significantly reduce future *Hylobius* populations, particularly if they are carried out within a few months of felling. For either of these techniques to be reasonably effective they must be undertaken prior to the first emergence of damaging adults from the stumps, which is normally a minimum of 14 months after felling on upland restock sites or 12 months in warmer lowland situations, in the current climate. However, it is almost certainly the case that some larvae will remain in roots, stems or branches left on site and, if so, these will still emerge as adult weevils to cause damage. Therefore, whilst mulching or de-stumping may be helpful techniques if they are practised for other reasons (e.g. on sites at high risk of *Heterobasidion annosum* (Fr.) Bref. infection, or to prepare a site for planting), they should not be relied on in isolation. These techniques can also pose a number of environmental risks and are not recommended in some situations, for example de-stumping should not be practised on sites classified as having a high risk of increased soil and water acidification, or where it will reduce site nutrient status or lead to an additional soil carbon loss (Moffat *et al.*, 2011).



### 3.4.3. Physical protection

Broadleaves or conifers protected by full tree shelters  $(1.2-1.8 \text{ m tall with no holes or porous mesh, and no gap between the bottom of the shelter and the ground) will suffer less damage than unprotected trees. The use of tall tree shelters purely for protection against$ *Hylobius*damage is likely to be prohibitively expensive, but if they are needed to prevent browsing mammals damaging trees, then any protection they provide against*Hylobius*is a bonus. Using tree shelters on areas greater than 1 ha is not normally economic, and they can cause instability in young conifers.

Other physical protection methods have been successfully used in Scandinavia, but trials in Britain have found them to be generally ineffective mainly, it is thought, due to the exceptionally high UK populations of *Hylobius* (Willoughby *et al.* 2020). Population levels in the UK and Ireland are often at least 7 times higher than those found on the most populous sites in Sweden (Willoughby *et al.* 2020), which is one reason that some non-chemical approaches that may be effective elsewhere in Europe are often not as successful here.

Physical protection products found to be effective in other countries, but less effective in the UK, include wax (Kvaae Wax<sup>®</sup>) and a polysaccharide stem coating (Flexcoat<sup>®</sup>), which are both flexible stem coatings applied before planting. A flexible stem coating system combining sand and glue (Conniflex<sup>®</sup>) needs further testing in UK conditions, as initial results from trials in this country were inconclusive. In addition, the following physical barriers were also tested and found to be generally ineffective in the UK unless *Hylobius* population levels are low:- custom made plastic weevil guards (MultiPro<sup>®</sup>, and Biosleeve<sup>®</sup>) fitted on site after planting; and lightweight plastic nets (WeeNets<sup>®</sup>) fitted around the root plug and lower stem of the tree at the nursery (Willoughby *et al.* 2020; Moore *et al.*, 2021).

Therefore, physical barriers such as MultiPro<sup>®</sup>, and to a lesser extent other products, may have a role in UK forestry in preventing unacceptable levels of *Hylobius* damage to newly planted trees, but only if the *Hylobius* Management Support System (Forest Research, 2021) or an alternative, reliable method of estimating population levels is used, and the onsite population is predicted to be low (i.e. untreated tree loss due to *Hylobius* browsing is expected to be < 25%). Even then, MultiPro<sup>®</sup> barriers should only be used on sheltered sites, and they need to be combined with the use of good-quality planting stock possessing large root collar diameters, an appropriate balance between the dry weight of their roots and shoots (Morgan, 1999) and with few side branches, which may limit their use in practice to vegetatively propagated material. Site preparation to create a weed and brash free area around the planted tree is also essential (Figure 4). In these circumstances, physical barriers, as part of an integrated approach, may be worth considering as a potentially suitable, if more expensive, substitute for the use of insecticides in the UK.



### 3.4.4. Insecticides

Unfortunately, in the UK and Ireland, even if a combination of non-chemical strategies are used it is often not possible to avoid the problem of *Hylobius* damage to young trees. In these circumstances, where there are no practical alternatives not involving excessive cost, the only option remaining to allow forest regeneration may be to treat young plants with insecticide. Used correctly, this can be a safe, effective, and economic means of protecting trees. Insecticides can be used before planting ('pre-planting pre-treatment'), during or immediately after planting ('post-planting'), possibly combined with further reapplications at intervals if weevil populations are high and persistent ('post-planting topup spraying'). Whilst this strategy normally provides very effective protection of seedlings, it has little effect on total populations of weevils on site and when used in isolation, if populations are very high, occasionally the insecticide protection can be overcome.

Currently, the only insecticides approved by the UK Health and Safety Directorate for use in forests to protect against *Hylobius* damage are alpha-cypermethrin, cypermethrin, and acetamiprid (the approval for chlorantraniliprole ended in December 2021). Alpha-cypermethrin and cypermethrin are non-systemic pyrethroid insecticides with high activity against *Hylobius*, but they are both potentially very toxic to the aquatic environment if misused, even if only very small quantities were to pollute watercourses.

Alpha-cypermethrin is only used to treat trees off-site in horticultural tree nurseries, before planting in the forest, and therefore presents a very low risk of damage to the wider environment. Consequently, use in this way is permitted on estates voluntarily certified to Forest Stewardship Council standards via the UK Woodland Assurance Standard (UKWAS, 2018; FSC 2019).

Cypermethrin had been the preferred insecticide for post-planting top-up sprays to protect young trees from high population levels of *Hylobius* for several years, but environmental concerns led to a collaborative research effort across the UK forest industry to search for less hazardous alternatives. The concerns around cypermethrin have been over: a) Its toxicity to aquatic life if misused; b) Its classification as a priority substance under the European Commission Water Framework Directive meaning emissions have to be progressively reduced, and; c) The fact that it is classified as 'highly hazardous – highly restricted' by the Forest Stewardship Council, the presumption being that it should not normally be used for post-planting top-up sprays on estates voluntarily certified to FSC standards via the UK Woodland Assurance Standard (UKWAS) where there are less hazardous alternatives available (UKWAS, 2018; FSC 2019).

As already noted, recent research found that physical protection products were only of use in certain circumstances in the UK. As part of this work, a wide range of repellents, and natural product insecticides such as garlic, azadirachtin derived from Neem oil, spinosad and pyrethrins were also tested and unfortunately found to be ineffective.



However, the synthetic insecticides acetamiprid and chlorantraniliprole were found to be highly effective in preventing *Hylobius* damage to young trees (Hardy *et al.*, 2020; Willoughby *et al.*, 2020; Moore *et al.*, 2021).

Acetamiprid is a systemic neonicotinoid insecticide that is widely used to control aphids and other damaging insects in agricultural and horticultural crops such as apples, tomatoes, potatoes and oilseed rape, as well as being used extensively in home garden insecticides. Acetamiprid is at least 500 times less toxic to aquatic life than cypermethrin, based on Predicted No Effect Concentrations (PNEC) of 500 ng/l for acetamiprid and 1 ng/l for cypermethrin (European Commission, 2004a; 2005), although when formulated as the pesticide product Gazelle SG<sup>®</sup> it is still classed as potentially very toxic to aquatic life if misused, due to its effect on the early life stages of aquatic insects - see Table 3 and the Appendix for further details. In recent years it has therefore increasingly been phased in as a replacement for alpha-cypermethrin for preplanting treatment, and cypermethrin for post-planting top-up sprays.

Chlorantraniliprole is a relatively low toxicity, selective, translaminar, synthetic diamide insecticide acting by contact and ingestion. It is widely used to control caterpillars on fruit and vegetable crops. It is not hazardous to mammals, and has a similar level of toxicity to bees as acetamiprid (EFSA, 2013) – see Table 3. In forestry, it only had approval for pre-planting treatment of trees, and this ended on December 31<sup>st</sup> 2021. This method of preventing *Hylobius* damage is therefore not covered in detail here. However, it is likely that new chlorantraniliprole products will become available for use in pre- and post-planting situations within the next few years.

## 3.4.4.1 Pre-planting pre-treatment with alpha-cypermethrin (Alpha C 6ED<sup>®</sup>) applied through an Electrodyn<sup>®</sup> spray booth

This is a specialised, fully enclosed application system located in industrial buildings in horticultural tree nurseries or depots. Insecticide is not used in a forest environment, but as part of the nursery plant production process to pre-treat trees before they reach the forest. The machinery generates electrostatically charged spray droplets that are attracted to the earthed transplants passing on a conveyor belt, enabling use of small volumes of insecticide (0.1 ml per tree of Alpha C 6ED<sup>®</sup>, equivalent to 0.006 g of alpha-cypermethrin per plant) allowing rapid drying, and providing consistent positioning of the band of insecticide onto the lower part of the stem and root collar of the tree only. The Electrodyn<sup>®</sup> system can only be used with specially formulated pesticides, in this case the forest nursery approved product Alpha C 6ED<sup>®</sup> (60 g/l alpha-cypermethrin; Techneat, 2007).

Treating plants in this way ensures a high level of operator protection, and allows plants to be transported soon after application without the need for a prolonged period of drying. Because pre-treated plants are delivered dry with the insecticide having been absorbed into the tree bark, as long as the trees are not put directly into watercourses, there is very little risk of any environmental contamination. Although the active



ingredient alpha-cypermethrin itself is toxic to bees, there is almost no risk of exposure to bees when it is used in this way (see the Appendix for further detail on the risk of exposure to bees from insecticide treated trees).

The product Alpha C 6ED<sup>®</sup> is harmful if swallowed or inhaled, so in addition to adopting protective measures when it is being applied using the Electrodyn<sup>®</sup> machinery, anyone handling or planting treated trees must wear the appropriate personal protective equipment, and vent planting bags for 1 hour on site after delivery to the forest to allow any build-up of the volatile naphtha solvent oil to dissipate (see FISA, 2013a+b). Accidental contamination of forest users by alpha-cypermethrin is highly unlikely as only the lower part of the stem of relatively small trees is treated. In addition, as the insecticide is dry it is unlikely to transfer through accidental contact, and it is also not hazardous in contact with the skin. A full review of the hazard characteristics of alphacypermethrin is given in European Commission (2004b). A summary of some of these characteristics, for comparison with other insecticides approved for use against Hylobius, is given in Table 3. The Alpha C 6ED<sup>®</sup> product label and approvals document (Health and Safety Executive, 2007; Techneat, 2007) are essential reading and give full details of application methods and other important information for the safe and effective use of the pesticide. The approval for Alpha C 6ED<sup>®</sup> ends on January 31<sup>st</sup> 2023. Currently, no other products are approved for use in the Electrodyn<sup>®</sup> machinery, and therefore after this date this particular method of treatment will no longer be available.

## 3.4.4.2 Pre-planting pre-treatment with acetamiprid (Gazelle SG $^{\rm (B)}$ ) applied via spray booth machinery

This treatment is applied using various different designs of purpose-built spray booth machinery in the nursery to generate a conventional aqueous spray delivering no more than 0.185 g of Gazelle SG<sup>®</sup> product (20% w/w acetamiprid; Certis, 2017) per plant, or 0.037 g acetamiprid active ingredient per plant. The spray booth provides controlled treatment with minimal operator exposure and the plants are left under cover to dry before despatch. Again, in addition to the usual requirement to protect operators when spraying, anyone handling or planting treated trees must wear the appropriate personal protective equipment (see FISA, 2013b). In contrast to trees pre-treated with alpha-cypermethrin via the Electrodyn<sup>®</sup>, plants treated with acetamiprid in spray booths can be cold stored for at least 3 months without harming the trees, and there is some evidence that doing so actually increases the efficacy of protection against *Hylobius* (Moore *et al.*, 2021).

As with the Electrodyn<sup>®</sup> system, plants are treated off-site in an industrial building in a horticultural tree nursery, and are delivered with the insecticide having been absorbed by the tree. Therefore, as long as the trees are not put directly into watercourses, and the plastic bags plants are delivered to site in are not placed in or near road drains or watercourses and removed from the site at the end of the day, there is very little risk of any harmful environmental contamination, or risk to forest users or bees. This has been



confirmed by a recent Forest Research study (Thomas *et al.*, in prep.) – see the Appendix for further details.

Details of the hazard characteristics of acetamiprid, based on the advice issued by the Health and Safety Executive, are given in the Appendix. A summary of some of these characteristics, for comparison with other insecticides approved for use against *Hylobius*, is given in Table 3. The Gazelle SG<sup>®</sup> product label and approvals documents (Health and Safety Executive 2011; 2012; 2016; 2017; Certis, 2017) are essential reading and give full details of application methods and other important information for the safe and effective use of this pesticide.

#### 3.4.4.3 Immediate post-planting spraying with cypermethrin or acetamiprid

If pre-treated trees are not used, it may be possible on a small scale to spray untreated trees immediately after planting with cypermethrin or acetamiprid (see below). This operation must be carried out as soon as stock is planted to avoid the high risk that the unprotected plants will be damaged or killed, but it should be noted that acetamiprid can only be applied between March to October. Both conifers and broadleaves can be treated. However, a cheaper and more effective approach, particularly on larger scale restock sites, is to use trees pre-treated in the nursery with acetamiprid or alpha-cypermethrin before planting. In addition, as noted above, nursery treatments can be applied in more controlled conditions, and the resulting risk to the wider environment from the insecticide application is considerably lower.

#### 3.4.4.4 Post-planting top-up spraying with cypermethrin or acetamiprid

Even if pre-treated trees are used, top-up spraying of cypermethrin or acetamiprid is sometimes needed in the autumn of the first year of planting, normally required in the spring of the second year after planting, and occasionally in the spring of the third year after planting, unless *Hylobius* population levels are low or if a fallow period strategy has been followed. Acetamiprid usually provides a greater duration of protection than alpha-cypermethrin or cypermethrin.

Typically, if insecticide pre-treated Sitka spruce (*Picea sitchensis* (Bong.) Carrière) with larger root collar diameters (e.g.  $1\frac{1}{2} + 1\frac{1}{2}$  stock) are used, by the autumn of the second year after planting trees are often large enough to resist attack. Smaller plants, or those of other species such as Douglas fir (*Pseudotsuga menziesii* (Mirbel) Franco), may need protecting for up to 4 years or more after planting. In all cases the risk is dependent on residual weevil population density on site which in turn depends on factors such as the length of fallow period adopted before planting. Young trees need to be monitored closely during the peak damage times (April – June, and August – October) and treated at the first sign of damage.

For some species and sites, if damage starts to occur, further treatment may be required in the spring and autumn of subsequent years. However, poor weather conditions, particularly rain or wind, may prevent a site being sprayed immediately damage is



recorded, which may mean that trees are damaged so severely that they die before they can be sprayed with insecticide.

Therefore, rather than simply reacting to damage when it occurs, it is recommended that the *Hylobius* Management Support system, or similar, is used to predict likely future browsing risk. This will help avoid damage by allowing the application of a well-timed top-up spray before the onset of browsing, and will reduce the likelihood of unnecessary prophylactic spraying taking place.

It is recommended that a marker dye such as a 2% solution of the food dye Dysol Turquoise (to give an equivalent of 0.8% active ingredient Acid Blue 9 in the final spray volume) is used to help target the top-up spray, minimise drift, and prevent unnecessary run-off (Brown *et al.*, 2003). Only trained operators can legally apply pesticides in the forest, and the legal requirements are summarised in the appropriate Pesticides Code of Practice (DEFRA, 2006; Scottish Executive, 2006). Normal good working practices for top-up spraying, including required personal protective equipment, are summarised in FISA (2013a).

Forester<sup>®</sup> (100g/l cypermethrin; Fargro, 2009) is approved for post-planting use in forests. It should be applied as a 10–20 ml spray of a 2% solution of Forester<sup>®</sup> in water (i.e. 20 ml of Forester<sup>®</sup> in 1 litre of water, equivalent to a 0.2% solution of cypermethrin active ingredient), using a handheld knapsack sprayer or spot gun or equivalent. This results in an equivalent of up to 0.4 ml of Forester<sup>®</sup> being applied per tree, or 0.04 g per tree of cypermethrin active ingredient. In conifers, feeding starts, and is most severe, in the upper parts of the roots and lower section of the main stem. Therefore, the spray should be carefully directed to cover the entire circumference of lower half of the stem, allowing the solution to flow down the stem and onto the root collar (Figure 5). Run-off into the soil, or drift to surrounding soil or vegetation should be minimised by the use of low spraying pressures (around 1 bar) and suitable nozzles, such as an adjustable cone nozzle. In broadleaves, *Hylobius* starts to feed at the main bud and progresses down the stem, so the upper part of the tree stem should be treated as well as the lower part. Three applications are permitted each year.

A full review of the hazard characteristics of cypermethrin is given in European Commission (2005). A summary of some of these characteristics, for comparison with other insecticides approved for use in protecting against *Hylobius*, is given in Table 3. The Forester<sup>®</sup> product label and approvals document (Health and Safety Executive, 2009; Fargro, 2009) are essential reading and give full details of application methods and other important information for the safe and effective use of this pesticide.

Gazelle SG<sup>®</sup> (20% w/w acetamiprid; Certis, 2017) is also approved for post-planting use in forests. It should be applied as a 10–20 ml spray of a 0.92% solution of Gazelle SG<sup>®</sup> in water (i.e. 9.25 g of Gazelle SG<sup>®</sup> in 1 litre of water, equivalent to a 0.184% solution of acetamiprid active ingredient), using a handheld knapsack sprayer or spot gun or equivalent fitted with a suitable nozzle such as an adjustable cone nozzle. This results in



an equivalent of up to 0.185 g of Gazelle SG<sup>®</sup>, or 0.037 g of active ingredient per tree. The spray should be carefully directed to cover the entire circumference of lower half of the stem, allowing the solution to flow down the stem and onto the root collar. In addition the foliage should be treated, as this allows the systemic insecticide to be absorbed and then translocated around the plant (Figure 5). Run-off into the soil, or drift to surrounding soil or vegetation, should be minimised by using low spraying pressures (around 1 bar) and suitable nozzles (such as an adjustable cone nozzle). Applications can only be made between March to October. A maximum of 416.25 g/ha of Gazelle SG<sup>®</sup> or 83.25 g of acetamiprid can be applied per hectare per application, and two applications are permitted each year. For each application this is an equivalent of treating 2250 trees at 0.185 g of Gazelle SG<sup>®</sup> per tree and gives the optimum levels of protection, but for higher initial planting densities the dose rates per tree need to be reduced. Minimum rates of 0.111 g of Gazelle SG<sup>®</sup> per tree are still moderately effective, which would allow up to 3750 trees per hectare to be treated, whilst staying within the maximum permitted dose per hectare. However, there are some indications that on the sites with the highest Hylobius population levels, the lower minimum rate of 0.111 g per tree may not be sufficient to prevent all damage (Willoughby et al., 2020). Therefore on the sites which are predicted to have the very highest Hylobius population levels, if the aim is to reduce the number of trees that need to be replaced (beaten up), it may be better to reduce initial planting densities as far as possible towards 2250 stems per hectare, to allow as close as possible to the maximum rate of 0.185 g of Gazelle SG<sup>®</sup> per tree to be used.

Details of the hazard characteristics of acetamiprid, based on the advice issued by the Health and Safety Executive, are given in the Appendix. A summary of some of these characteristics, for comparison with other insecticides approved for use against *Hylobius*, is given in Table 3. The Gazelle SG<sup>®</sup> product label and approvals documents (Health and Safety Executive 2011; 2012; 2016; 2017; Certis, 2017) are essential reading and give full details of application methods and other important information for the safe and effective use of this pesticide.



Table 2. A summary of the relative costs, efficacy and potential risks of the main remedial control measures available for protecting trees from damage by *Hylobius abietis* 

Method	Approximate cost per treated ba	Approximate total cost per ha for	Efficacy	Summary of potential environmental risks	Comments						
	per operation <sup>1</sup>	satisfactory restocking <sup>1,2</sup>									
Non-chemical methods											
Cultivation to expose mineral soil	£200 - 600 <sup>3</sup>	Not possible	Can give some benefit, but unlikely to reduce damage to an acceptable level by itself.	Soil erosion, water sedimentation, nutrient leaching, soil carbon loss, especially on organic soils, destruction of soil fauna, disruption of ground nesting birds and archaeology, and possible atmospheric pollution (especially if machinery is poorly maintained) can all result from inappropriate cultivation if guidelines are not followed.	Mounding or scraping of the soil at the planting position is necessary. This technique will result in the greatest reduction in weevil damage when used on sites with a high density of woody weeds. Planting should be in the centre of the cleared area of mineral soil.						
Stump removal	Very variable, but can be in the region of £450–650	Not possible	Can help to reduce damage but may not be wholly effective by itself.	Soil and habitat disturbance, potential soil compaction. On steep sites, potential soil erosion and water sedimentation. Likely increase in decomposition of soil organic matter leading to a reduction of soil carbon stocks. Possible atmospheric pollution from machine use, especially if it is poorly maintained. Removal of stumps from site will result in some loss of nutrients and an increased risk of soil acidification.	De-stumping will be more effective if carried out within a few months of felling. However, some larvae are likely to remain in any roots, stems or branches left on site. Removal of stumps is only worth considering where there is a ready market for the resulting biomass, and where sites are flat and relatively easy to de-stump with a low risk of negative impacts, or there are other particular benefits such as disease control. Stumps should not be windrowed, because <i>Hylobius</i> may still emerge from them. De-stumped ground is ideal for machine- planting the next rotation of trees.						
Mulching	£500–2000 <sup>3</sup>	Not possible	Can help to reduce damage but may not be wholly effective by itself.	Severe soil and habitat disturbance, although probably somewhat less than for stump removal. Risk of possible atmospheric pollution from machine use, especially if it is poorly maintained, damage to soil and overstorey stems from flailing and any subsequent burning, and atmospheric pollution from burning.	Mulching is likely to be somewhat more effective than de-stumping, if it also destroys roots and branches. Mulching will be more effective if carried out within a few months of felling. However, some larvae are likely to remain in any roots, stems or branches left un- mulched on site. It is probably only worth considering if mulching is planned anyway to prepare site prior to planting, and even then careful monitoring of subsequent insect emergence and damage is required. The site may need to be raked and burnt or scarified after mulching to allow planting. Mulching is only worth considering where sites are flat, well drained and relatively easy to work with a low risk of negative impacts.						
Tree shelters	£4250 (£1.70 per 1.2 m tall shelter) <sup>3</sup> £5500 (£2.20 per 1.8 m tall shelter) <sup>3</sup>	£4250 £5500	Potentially effective, but only if shelters are properly fitted, regularly inspected and maintained.	Unless fully biodegradable or removed, tree shelters will form a source of plastic / petrochemical pollution at the end of their life. Many conifers are not suited to growth in tall tree shelters as they can become unstable when they emerge from the tops of the shelter.	Not economic for protection from <i>Hylobius</i> alone, but if being used anyway to protect trees from browsing mammals, their use may be a viable option. The shelters need to be removed after trees are established.						



### Management of Hylobius abietis

Proprietary physical barriers, e.g. MultiPro <sup>®</sup> , Biosleeve <sup>®</sup> , WeeNet <sup>®</sup>	Limited trials suggest a cost (materials; fitting and extra planting cost) of around £300 - £1100 ha <sup>-1</sup> , depending on the type of guard fitted, and whether or not plants are supplied with guards pre- fitted. This does not include maintenance costs which could be considerable, depending on design		In the UK, only effective on sites with low <i>Hylobius</i> population levels.	Unless fully biodegradable or removed, they will form a source of plastic / petrochemical pollution at the end of their life.	Physical barrier sleeves such as MultiPro <sup>®</sup> , Biosleeve <sup>®</sup> , and to a lesser extent WeeNets <sup>®</sup> c be effective but only if used in conjunction with the Hylobius Management Support System or alternative, reliable method of estimating population levels, and the onsite population is predicted to be low (i.e. < 25% untreated tree loss). In the UK, they are not generally sufficiently effective by themselves on sites w medium or high Hylobius population levels. Barriers should be combined with the use of good quality well balanced planting stock with large root collar diameters and short side branches (i.e. ideally vegetatively propagated material), and with suitable site preparation t create a weed and brash free site around the planted tree. The site should be level, and sheltered from strong winds. The site should I regularly inspected and the barriers maintain to ensure they remain firmly pushed into the ground and that no bridging from surroundin brash has occurred. Most designs of barrier a designed to split after the main risk from dama is over and in theory may not need to be removed to allow continued tree growth. Nor porous barriers may make subsequent insectic spraying, if required, difficult unless the guar are removed. To date most flexible stem coatings have bee found to be ineffective in the UK.		
Nematodes: Steinernema carpocapsae	Approximate cost of £670 per ha per year during the 3-4 treatment years.		Can be effective in allowing trees to establish on some sites.	Insect parasitic nematodes are naturally occurring and have no known adverse impacts on non- target organisms. Atmospheric pollution may be possible from application machinery, particularly if it is poorly maintained. Soil compaction is possible from machine trafficking.	If used in conjunction with the <i>Hylobius</i> Management Support System, on consecutive clearfell sites within large forest blocks, may over 3-4 years reduce population levels sufficiently such that no further remedial control measures are required (i.e. insecticide use is not required). Not suitable for small scale, sporadic or isolated clearfells, or on steep sites or where the high volumes of water required cannot be sourced or transported.		
Chemical methods							
Insecticides, general			Can be very effective, even on sites with high <i>Hylobius</i> population pressure.	If misused, all insecticides present a risk to operator health, risk of soil and water pollution, potential risk of poisoning of wildlife and damage to non-target vegetation and insects. Risks are reduced by spot treatment in the forest or use under highly controlled conditions in off-site horticultural tree nurseries. No long- term harmful effects have been detected at planting sites.			
Pre-plant pre-treatment of trees with alpha- cypermethrin applied via Electrodyn <sup>®</sup> spray booth	£130	Complete protection rarely possible, usually needs to be combined with top-up spraying	Provides very good protection for first season, but dependent on weather and pest population	Used only in highly controlled conditions in off-site horticultural tree nurseries, where if not adequately controlled the insecticide could be harmful if swallowed or by inhalation. Once the insecticide has dried on stems, there is believed to be a low risk to the wider environment when treated trees are planted in the forest, but planters still need to wear PPE.	Only suitable for bare-rooted stock treated before planting. The plants usually require additional top-up treatment(s) in the second and sometimes the third year after planting, particularly when insect populations are high and if small sized plants are used <sup>4</sup> .		



### Management of Hylobius abietis

Pre-plant pre-treatment of trees with acetamiprid	£170 - £225	Complete protection rarely possible, usually needs to be combined with top-up spraying	Provides very good protection for first season, but dependent on weather and pest population.	Used in highly controlled conditions in off-site horticultural tree nurseries, where if not adequately controlled the insecticide could be harmful if swallowed. Once the insecticide has dried on stems, there is believed to be a low risk to the wider environment when treated trees are planted in the forest, but planters still need to wear PPE.	The plants usually require additional top-up treatment(s) in the second and sometimes the third year after planting, particularly when insect populations are high, and if small sized plants are used <sup>4</sup> .
Post-plant top-up spraying with cypermethrin	£140	£420 - £1120	Provides good protection for around 8-10 weeks after application, depending on population levels.	With proper controls in place cypermethrin can be used safely in the forest, but cypermethrin is a broad spectrum insecticide and if misused many non-target insects could be killed. It is harmful if swallowed, may cause an allergic skin reaction, and very toxic to aquatic life with long lasting effects. Once the insecticide has dried on stems, there is believed to be a low risk to the wider environment.	Treatment must take place during dry weather and all normal controls and good working practices when using pesticides must be in place. The deposit becomes rain fast once dry. Applications should be timed such that a minimum rain free period of at least 1 hour, and preferably 6 hours, occurs after spraying. Dyes may be added to the solution to check coverage. Plants typically need treatment for at least 2 years after planting with 1-2 treatments a year, depending on population levels. Population and likely damage due to <i>Hylobius</i> can be predicted by using the <i>Hylobius</i> Management Support System, which avoids the need for unnecessary prophylactic sprays or waiting for damage to occur before treating. If combined with the use of pre-treated trees, top-up spraying may only be required (twice) for the second year after planting (i.e. total cost of protecting trees to establishment of around £410 /ha). For some species and sites top-up spraying may be required for 4 years or more after planting <sup>4</sup> . Individual treatment costs, efficacy and environmental risks are identical for immediate post-planting spraying of cypermethrin.
Post-plant top-up spraying with acetamiprid	£160	£320 - £1280	Provides as good as or better protection than cypermethrin due to its systemic activity, i.e. effective for at least around 8-10 weeks after application, depending on population levels.	With proper controls in place acetamiprid can be used safely in the forest, and compared to cypermethrin it is more selective, less hazardous to the aquatic environment and less hazardous to bees. However, if misused acetamiprid is harmful if swallowed, and very toxic to aquatic life with long lasting effects. Once the insecticide has dried on stems, there is believed to be a low risk to the wider environment.	Treatment must take place during dry weather and all normal controls and good working practices when using pesticides must be in place. The deposit becomes rain fast once dry. Applications should be timed such that a minimum rain free period of at least 1 hour, and preferably 6 hours, occurs after spraying. Dyes may be added to the solution to check coverage. Plants typically need treatment for at least 2 years after planting with 1-2 treatments a year, depending on population levels. Population and likely damage due to <i>Hylobius</i> can be predicted by using the <i>Hylobius</i> Management Support System, which avoids the need for unnecessary prophylactic sprays or waiting for damage to occur before treating. If combined with the use of pre-treated trees, top-up spraying may only be required (once) for the second year after planting (i.e. total cost of protecting trees to establishment of around £330 /ha). For some species and sites top-up spraying may be required for 4 years or more after planting <sup>4</sup> . Individual treatment costs, efficacy and environmental risks are identical for immediate post-planting spraying of acetamiprid.

#### Notes

1 Costs are approximate and are included for comparison purposes only, and may not reflect the actual cost achievable in particular circumstances and locations. The cost for individual insecticides includes chemical plus application costs, assuming 2500 trees being treated per hectare.



- 2 The total cost figure includes any repeat treatment that may be required to achieve satisfactory restocking using only this single method. In reality, a combination of different techniques is often required.
- 3 This operation may be being undertaken in any case for its silvicultural benefits, in which case there is effectively no marginal cost for any additional *Hylobius* protection obtained.
- 4 The number of years for which the plants will require protection will also depend on factors such as the length of any pre-planting fallow period imposed and the size of the *Hylobius* population.

#### Figure 4. Tree planted on a cultivated, weed-free mound, protected with a Multipro® guard





#### Table 3. A comparison of some of the hazard characteristics of insecticides used for protecting trees from damage by Hylobius abietis

Pesticide	Mode of action	Rate applied active ingredient (kg ha <sup>.,</sup> )	Rate applied active ingredient (mg m <sup>2</sup> )	Active ingredient toxicity to mammals (rats): oral (LD50, mg kg <sup>4</sup> ) <sup>1</sup>	Active ingredient toxicity to mammals (rats): contact (LD50, mg kg <sup>4</sup> ) <sup>1</sup>	Active ingredient toxicity to mammals (rats): oral NOEL (mg kg- daily by body weight) (duration) <sup>2</sup>	Formulation toxicity to mammals (rats): oral (LD50, mg kg <sup>-1</sup> ) <sup>-1</sup>	Formulation toxicity to mammals (rats): contact (LD50, mg kg <sup>-1</sup> ) <sup>-1</sup>	Hazard classification of product formulations	Toxicity to invertebrates (bees): contact (LD <sub>50</sub> μg per bee) <sup>1</sup>	Hazard classification of product formulation: potential risk to aquatic life	FSC 'highly hazardous' list status	Persistence in soil DT 50 (days) <sup>3</sup>
Alpha- cypermethrin	Insecticide	0.02	2	57	>2 000	> 1.5	629	2 000	Harmful by inhalation or if swallowed, may cause lung damage if swallowed Skin sensitizer.	<b>Not toxic</b> <sup>4</sup> 0.059	Very toxic <sup>7</sup> to aquatic life with long lasting effects	'Highly hazardous – highly restricted'	91
Cypermethrin	Insecticide	0.05-0.1	5-10	250-4 150	>4 920	5 (2 years)	300-2 000	>2 000	Harmful if swallowed May cause an allergic skin reaction.	Not toxic <sup>4</sup> 0.02	Very toxic <sup>7</sup> to aquatic life with long lasting effects	'Highly hazardous – highly restricted'	35
Acetamiprid	Insecticide	0.08	8	146	>2 000	7.1 (2 years)	1 065	>2 000	Harmful if swallowed	Not toxic <sup>6</sup> 8.1	Very toxic <sup>7</sup> to aquatic life with long lasting effects	'Highly hazardous – restricted'	1-5
Chlorantranili prole	Insecticide	0.035	3.5	>5 000	>5 000	158 (18 months)	>5 000	>5 000	Not hazardous	Not toxic <sup>6</sup> >4	Very toxic to aquatic life with long lasting effects	'Highly hazardous – restricted'	60-365

#### Notes

This table is a collation based on published information from MacBean (2012), European Commission review reports, individual product material safety data sheets and product labels. No testing has taken place by Forest Research. The table is intended as a guide to the relative characteristics of different pesticides for general comparison purposes only, to aid in the selection of the least hazardous pesticide option, if there are no viable non-chemical alternatives. Different approved products with the same active ingredients may have different characteristics. Refer to the product label and material safety data sheet for specific products for definitive information on hazards and safety before making and final decision on pesticide selection, and before making any application. Note that just because a product is hazardous does not mean it is unsafe to use – as long as the risk of the hazard materialising is minimized, through adopting good forestry practices including, for example, following the instructions on product labels, employing trained operators, and using adequate PPE and no-spray buffer zones, then a potentially hazardous substance can still be used safely.

- 1 LD<sub>50</sub> is the lethal dose of the substance in milligrams of active ingredient for each kilogram of bodyweight of the target that, if ingested, is likely to kill 50% of the population. 1 mg = 1 milligram = 0.001 grams; 1  $\mu$ g = 1 microgram = 0.000,001 grams; 1 mg = 1 nanogram = 0.000,000,001 g. For comparison, caffeine has an LD<sub>50</sub> of 367 mg kg<sup>-1</sup> bodyweight (Adamson, 2016).
- 2 NOEL is the No Observable Effect Level, the amount that can consumed daily for each kilogram of body weight of the consumer with no observable effect.

3 DT<sub>50</sub> is the half-life for the substance in soil, the time taken in days for 50% of the substance to break down. It gives an approximate estimate only of likely persistence in soil. Actual breakdown rates will depend on soil type and weather conditions. Breakdown in water and plants is often quicker. Often a substance may be bound up and effectively safely immobilised in soil very rapidly, even though actual decomposition may take many days.

4 Although the active ingredient itself can clearly be highly toxic to invertebrates, the UK Health and Safety Executive have judged that the product formulation will not be harmful to bees when used as directed.

5 FSC pesticide policy does not apply to plants pre-treatment in off-site horticultural tree nurseries.

6 Although the active ingredient itself can clearly be toxic to invertebrates, the UK Health and Safety Executive have judged that the product formulation will not be harmful to bees when used as directed.

7 Acetamiprid is around 500 times less toxic to aquatic life than cypermethrin although the products themselves have similar broad hazard classifications – see the Appendix for details.



## 4. Conclusions

*Hylobius abietis* can cause catastrophic damage to newly planted trees, effectively preventing regeneration within recently felled areas of conifer forest. Forest Research recommend that an integrated approach to the management of this pest is adopted, firstly by understanding its life cycle and likely impacts, and then by considering the full range of potential approaches available to prevent it damaging young trees, if necessary by using different techniques in combination with each other. Although research into non-chemical approaches is ongoing, currently on many sites in the UK and Ireland it is likely that insecticides will still need to be used as part of the integrated management of *Hylobius*. Where insecticide use is unavoidable, pre-treatment of young trees in an off-site horticultural tree nursery with alpha-cypermethrin or acetamiprid, possibly combined with later post-planting top-up sprays in the forest of those trees with acetamiprid, when used as described on the Health and Safety Executive approved pesticide product labels, and when combined with the additional precautionary measures in place as described in this Report, should not pose any unacceptable risk to consumers, operators, bystanders, neighbours or the wider environment.

## 5. References

Adamson, R.H. (2016). The acute lethal dose 50 (LD<sub>50</sub>) of caffeine in albino rats. *Regulatory Toxicology and Pharmacology* **80**, 274-276. <u>https://doi.org/10.1016/j.yrtph.2016.07.011</u>

Brixey, J.M., Moore, R., and Milner, A.D. (2006). Effect of entomopathogenic nematode (*Steinernema carpocapsae* Weiser) application technique on the efficacy and distribution of infection of immature stages of the large pine weevil (*Hylobius abietis* L.) in stumps of Sitka spruce (*Picea sitchensis* Carr.) created at different times. *Forest Ecology and Management* **226**, 161-172. <u>https://doi.org/10.1016/j.foreco.2006.01.044</u>

Brown, A., Willoughby, I., Clay, D.V., Moore, R and Dixon, F.L. (2003). The use of dye markers as a method of reducing herbicide use and potential environmental impacts. *Forestry* **76** (4), 371-384. <u>https://doi.org/10.1093/forestry/76.4.371</u>

Certis (2017). Gazelle SG® product label. www.certiseurope.co.uk

DEFRA (2006). *Pesticides – Code of practice for using plant protection products*. DEFRA, London. <u>www.pesticides.gov.uk/safe\_use.asp?id=64</u>

EFSA (2013). Conclusion on the peer review of the pesticide risk assessment of the active substance chlorantraniliprole. *European Food Safety Authority Journal* **11** (6), 3143. <u>https://doi.org/10.2903/j.efsa.2013.3143</u>



EFSA (2016). Peer review of the pesticide risk assessment of the active substance acetamiprid. *European Food Safety Authority Journal* **14** (11), 4610. https://doi/10.2903/j.efsa.2016.4610

European Commission (1998). Council Directive 98/83/EC of 3 November 1998 on the quality of water intended for human consumption. *Official Journal of the European Union* **L 330**, 0032 – 0054.<u>www.ec.europa.eu</u>

European Commission (2001). Guidance for the setting of an acute reference dose (ARfD). *European Commission document 7199/VI/99 rev.5*. https://ec.europa.eu/food/plant/pesticides/approval\_active\_substances/guidance\_docum\_ents\_en

European Commission (2004a). Review report for the active substance acetamiprid. *European Commission working document SANCO/1392/2001 – final*. <u>www.</u> <u>ec.europa.eu/food/plant/pesticides\_en</u>

European Commission (2004b). Review report for the active substance alphacypermethrin. *European Commission document SANCO/4335/2000.* www.ec.europa.eu/food/plant/pesticides\_en

European Commission (2005). Review report for the active substance cypermethrin. *European Commission document SANCO/4333/2000*. <u>www.ec.europa.eu/food/plant/pesticides\_en</u>

European Commission (2015). Development of the first Watch List under the Environmental Quality Standards Directive. *European Commission Joint Research Centre Technical Report EUR 27142 EN.* <u>www.ec.europa.eu/jrc/en/publication/eur-scientific-</u> <u>and-technical-research-reports/development-first-watch-list-under-environmental-</u> <u>quality-standards-directive</u>

European Commission (2016). Acetamiprid. List of information, tests and studies which are considered as relied upon by the RMS for the evaluation with a view to the renewal of the active substance. Rapporteur Member State: The Netherlands. European Commission. www.ec.europa.eu/food/plants/pesticides\_en

European Commission (2017a). Final review report for the active substance acetamiprid. *European Commission working document SANCO/10502/2017 – Rev 4.* <u>www.ec.europa.eu/food/plant/pesticides\_en</u>

European Commission (2017b). Commission Regulation (EU) 2017/626 amending Annexes II and III to Regulation (EC) No 396/2005 of the European Parliament and of the Council as regards maximum residue levels for acetamiprid, cyantraniliprole, cypermethrin, cyprodinil, difenoconazole, ethephon, fluopyram, flutriafol, fluxapyroxad, imazapic, imazapyr, lambda-cyhalothrin, mesotrione, profenofos, propiconazole,



pyrimethanil, spirotetramat, tebuconazole, triazophos and trifloxystrobin in or on certain products. *Official Journal of the European Union* **L 96/1**. <u>www.ec.europa.eu</u>

European Commission (2021). *Approval of active substances* (webpage). The European Commission. <u>www.ec.europa.eu/food/plant/pesticides/approval active substances en</u> (accessed 30/11/21).

Fargro (2009). Forester product label. www.fargro.co.uk

FERA (2017). *Pesticide usage survey statistics*. FERA Sciences Ltd. <u>https://secure.fera.defra.gov.uk/pusstats</u> .

FISA (2013a). Application of pesticides by hand-held equipment. *Forest Industry Safety Accord Safety Guide* **202**. Forest Industry Safety Accord. <u>www.ukfisa.com</u>

FISA (2013b). Planting. *Forest Industry Safety Accord Safety Guide* **103**. Forest Industry Safety Accord. <u>www.ukfisa.com</u>

Forestry Commission (2016). *Forestry facts and figures 2016*. Forestry Commission, Edinburgh. <u>www.forestry.gov.uk/statistics</u>

Forestry Commission (2017). *The UK Forestry Standard.* Forestry Commission, Edinburgh. 232 pp. <u>www.forestresearch.gov.uk</u>

Forest Research (2021). *Hylobius management support system* (web page). Forest Research. <u>www.forestresearch.gov.uk/tools-and-resources/fthr/tree-health-and-protection-services/hylobius-management-support-system</u> (accessed 30/11/21).

FSC (2019). *Forest Stewardship Council Pesticides Policy*. FSC-POL-30-001 V3-0 EN. The Forest Stewardship Council, Bonn. <u>www.fsc.org</u>

Hamilton, G.J. and Christie, J.I. (1971). Forest management tables (metric). *Forestry Commission Booklet* **34**. HMSO, London. <u>www.forestresearch.gov.uk</u>

Hardy, C., Sayyed, I., Dunbar Leslie A., Dittrich A.D.K. (2020). Effectiveness of insecticides, physical barriers and size of planting stock against damage by the pine weevil (*Hylobius abietis*). *Crop Protection* **137**: 105307. https://doi.org/10.1016/j.cropro.2020.105307

Health and Safety Executive (2007). Authorisation for the plant protection product Alpha C 6ED, MAPP Number 13611. *Authorisation number 02228 of 2007*. <u>www.hse.gov.uk/pesticides</u>

Health and Safety Executive (2009). Authorisation for the plant protection product Forester, MAPP Number 13164. *Authorisation number 0261 of 2009*. <u>www.hse.gov.uk/pesticides</u>



Health and Safety Executive (2011). Extension of authorisation for a minor use of the plant protection product Gazelle SG. *Extension of authorisation number 1313 of 2011*. www.hse.gov.uk/pesticides

Health and Safety Executive (2012). Extension of authorisation for a minor use of the plant protection product Gazelle SG. *Extension of authorisation number 1068 of 2012*. <u>www.hse.gov.uk/pesticides</u>

Health and Safety Executive (2016). Extension of authorisation for a minor use of the plant protection product Gazelle SG. *Extension of authorisation number 0084 of 2016*. <u>www.hse.gov.uk/pesticides</u>

Health and Safety Executive (2017). Authorisation for the plant protection product Gazelle SG, MAPP 13725. *Authorisation number 0260 of 2017*. <u>www.hse.gov.uk/pesticides</u>

Health and Safety Executive (2021). *Pesticides: The basics* (webpage). The Health and Safety Executive. <u>https://www.hse.gov.uk/pesticides/about-pesticides.htm</u> (accessed 30/11/21).

Loos, R., Marinov, M., Sanseverino, I., Napierska, D. and Lettieri, T. (2018). Review of the 1st Watch List under the WFD and recommendations for the 2nd Watch List. *JRC Technical Report*. <u>www.ec.europa.eu/jrc/en/publication/review-1st-watch-list-under-water-framework-directive-and-recommendations-2nd-watch-list</u>

Lundin, O., Rundlöf, M., Smith, H.G., Fries, I. and Bommarco, R. (2015). Neonicotinoid insecticides and their impacts on bees: A systematic review of research approaches and identification of knowledge gaps. *PLoS ONE* **10** (8): e0136928. https://doi.org/10.1371/journal.pone.0136928.g002

MacBean, C. (Ed.) (2012). *The pesticide manual, sixteenth edition*. BCPC, Alton. https://www.bcpc.org

Mason, W.L., Kerr, G., Pommerening, A., Edwards, C., Hale, S.E., Ireland, D., Johnson, D. and Moore, R. (2004). Continuous cover forestry in British conifer forests. *Forest Research Annual Report and Accounts 2003-04.* The Stationary Office, Edinburgh. p 38-53. <u>www.forestresearch.gov.uk</u>

McCarthy, N., Bentsen, N.S., Willoughby, I. and Balandier, P. (2011). The state of forest vegetation management in Europe in the 21st century. *European Journal of Forest Research* **130**, 7-16.

Moffat, A.J., Nisbet, T.R. and Nicoll, B. (2011). Environmental effects of stump and root harvesting. *Forestry Commission Research Note* **9**. Forestry Commission, Edinburgh. <u>www.forestresearch.gov.uk</u>



Moore, R. (2004). Managing the threat to restocking posed by the large pine weevil, *Hylobius abietis*: The importance of time of felling of spruce stands. *Forestry Commission Information Note* **61**. Forestry Commission, Edinburgh. <u>www.forestresearch.gov.uk</u>

Moore, R., Willoughby, I.H., Andrew J. Moffat, A.J. and Forster, J. (2021). Acetamiprid, chlorantraniliprole, and in some situations the physical barriers MultiPro<sup>®</sup> or Kvaae<sup>®</sup> wax, can be alternatives to traditional synthetic pyrethroid insecticides for the protection of young conifers from damage by the large pine weevil *Hylobius abietis* L. *Scandinavian Journal of Forest Research* **36** (4), 230-248.

https://doi.org/10.1080/02827581.2021.1906313

Morgan, J. L. (1999). Forest tree seedlings – best practice in supply treatment and planting. *Forestry Commission Bulletin* **121**. Forestry Commission, Edinburgh. <u>www.forestresearch.gov.uk</u>

Paterson, D.B. (1990). The potential to apply different silvicultural systems to upland forests, predominantly of Sitka spruce. In *Silvicultural Systems,* P. Gordon (Ed.). Institute of Chartered Foresters, Edinburgh, 120-138.

PEFC (2020). *PEFC UK certification scheme for sustainable forest management*. PEFC UK Ltd. <u>www.pefc.co.uk</u>

Scottish Executive (2006). *Pesticides – code of practice for using plant protection products in Scotland*. Scottish Executive, Edinburgh. www.scotland.gov.uk/Publications/2006/12/19110050/20

SEPA (2017). *The water environment (controlled activities) Scotland) regulations 2011 (as amended), a practical guide, version 7.6.* Scottish Environmental Protection Agency. <u>www.sepa.org.uk/regulations/water</u>

Techneat (2007). *Alpha C 6ED<sup>®</sup> product label*. <u>www.forestry.gov.uk/forestry/infd-5tug8w</u>

Thorsen, A.A., Mattsson, S., Weslien, J. (2001). Influence of stem diameter on the survival and growth of containerized Norway spruce seedlings attacked by pine weevils (*Hylobius* spp.). *Scandinavian Journal of Forest Research* **16** (1), 54-66. <u>https://doi.org/10.1080/028275801300004415</u>

Thomas, H., Nisbet, T.R. and Willoughby, I.H. (In prep.). Acetamiprid used to prevent *Hylobius abietis* damaging young trees poses a very low risk of causing water contamination provided good forestry practices are followed. In preparation for *Forestry*.

UKWAS (2018). *The UK woodland assurance standard: fourth edition, version 4*. UKWAS. <u>www.ukwas.org.uk</u>

Wainhouse, D., Brough, S. and Greenacre, B. (2007) *Managing the pine weevil on lowland pine. Forestry Commission Practice Note* **14**. Forestry Commission, Edinburgh, 12 pp. <u>www.forestresearch.gov.uk</u>



Willoughby, I., Evans, H., Gibbs, J., Pepper, H., Gregory, S., Dewar, J., Pratt, J., McKay, H. (2004). Reducing pesticide use in forestry. *Forestry Commission Practice Guide* **15**. Forestry Commission, Edinburgh. 140 pp. <u>www.forestresearch.gov.uk</u>

Willoughby, I., Balandier, P., Bentsen, N.S., McCarthy, N. and Claridge, J. (Eds.) (2009). *Forest vegetation management in Europe: current practice and future requirements*. COST Office, Brussels. 156 pp. <u>www.forestresearch.gov.uk</u>

Willoughby, I.H., Moore. R., Moffat, A.J, Forster, J., Sayyed, I. and Leslie, K. (2020). Are there viable chemical and non-chemical alternatives to the use of conventional insecticides for the protection of young trees from damage by the large pine weevil *Hylobius abietis* L. in UK forestry? *Forestry* **93** (5), 694–712. https://doi.org/10.1093/forestry/cpaa013

#### Figure 5. Post-planting top-up spraying of individual trees with insecticide and dye marker





## 6. Appendix - Acetamiprid safety

### 6.1. Introduction

Acetamiprid is a systemic, neonicotinoid insecticide that is widely used to control aphids and other damaging insects in agricultural and horticultural crops such as apples, tomatoes, potatoes and oilseed rape, as well as being used extensively in home garden insecticides.

Acetamiprid use in forestry is very different to conventional agricultural insecticide spraying. Instead of mechanised boom sprayers on tractors applying a chemical to entire fields, forest use of acetamiprid involves carefully directing a maximum of 20 millilitres of the dilute insecticide formulation (an equivalent of 0.185 g per tree of Gazelle SG<sup>®</sup>, or 0.037 g per tree of acetamiprid) via handheld applicators as a targeted spray, minimising run-off, onto the vulnerable part of the stem and foliage of each young tree (which are usually between 15 centimetres to 1 metre in height) without overspraying the surrounding soil or vegetation. Unlike arable or horticultural crops, insecticide applications in forestry are normally only made once or twice a year for the first 2-3 years of a typical forest rotation of 50-80 years.

Pesticide surveys have estimated that a total of 425,733 kg of insecticide active ingredient (a.i.) was used in agricultural and horticultural situations in GB in 2015, and this included 308 kg of acetamiprid (FERA, 2017). There are no equivalent surveys of insecticide use in forestry by which to make direct comparisons, but likely future annual usage can be very roughly estimated based on the total area of forests that are restocked with coniferous species, which in 2015 was reported to be 9,900 ha (Forestry Commission, 2016). On average at least 15% of the area of any restocking site is occupied by roads, rides, and watercourses (Hamilton and Christie, 1971), and so would never be treated. Of the remaining 8,400 ha, most sites would not be treated either because Hylobius populations did not warrant it, or because other methods of avoiding the problem or other non-chemical remedial control methods could be adopted. Based on typical past insecticide usage in the state forest sector, including sites treated in multiple years, it is unlikely that more than 40% of the total restocking area would ever be treated, giving a maximum GB area treated of very roughly around 3,500 ha per year, equating to an estimated realistic maximum annual use in GB forestry at current restocking levels of around 300 kg a.i. acetamiprid (this excludes the amount of acetamiprid used to treat trees in horticultural nurseries that are then subsequently planted out in the forest). As very few if any other insecticides are likely to be used, the total amount of active ingredient of insecticides applied in GB forests each year is also unlikely to exceed 300 kg.

In judging whether or not it is safe to use an insecticide such as acetamiprid, Forest Research, and the wider forest industry, relies on the expert advice and guidance



provided by the Chemicals Regulation Division of the UK Health and Safety Executive, whose primary focus is the protection of human health and the environment (Health and Safety Executive, 2021).

### 6.2. The approvals process for acetamiprid

Acetamiprid has been subject to international peer review of toxicology and environmental impact by the (then) 28 member states of the European Union (European Commission, 2021), which included an assessment of over 400 studies reporting on the insecticide's toxicology and environmental impacts (European Commission, 2016). The chemical has been accepted as safe to use in Europe (European Commission, 2004a; 2017a). In 2015, acetamiprid was placed on a 'Watch List' under the Environmental Quality Standards Directive for the purpose of collecting more high quality water monitoring data to help determine whether any further risk reduction measures are required (European Commission, 2015). The Watch List was reviewed in 2018. While the very low level of acetamiprid detected in sampled river waters fulfilled the criteria for its removal from the list, it was decided to retain the chemical on the Watch List as part of a general assessment of neonicotinoid concentrations in water (Loos *et al.*, 2018).

In the UK, pesticide use is subject to stringent government controls through legislation such as the Control of Pesticides Regulations, the UK implementation of the Sustainable Use Directive and the Plant Protection Products Directive, and the Control of Substances Hazardous to Health Regulations. Regulatory control is enforced by the Health and Safety Executive (Health and Safety Executive, 2021). After expert scientific scrutiny of all available safety data, the Health and Safety Executive have judged that acetamiprid, and the product Gazelle SG<sup>®</sup> (20% w/w acetamiprid; Certis, 2017), do not pose an unacceptable risk to consumers, operators, bystanders or the wider environment if used according to the conditions of use they have specified on the approved product label, and hence have granted approval for its use in agricultural, horticultural and home garden situations (Certis, 2017; Health and Safety Executive, 2017).

Because research showed acetamiprid to be as effective as cypermethrin in protecting young transplants from damage by *Hylobius* (Willoughby *et al.* (2020); Moore *et al.*, 2021), whilst being at least 500 times less toxic to aquatic life than cypermethrin (based on Predicted No Effect Concentrations (PNEC) of 500 ng/l for acetamiprid and 1 ng/l for cypermethrin; where 1 ng = 0.00000001 g) (European Commission, 2004a; 2005), the forest industry applied to the Chemicals Regulation Division of the Health and Safety Executive for an Extension of Authorisation for Minor Use (often abbreviated to 'EAMU') to use Gazelle SG<sup>®</sup> (containing acetamiprid) in forests. After reviewing all of the safety data, the Health and Safety Executive judged that this new use did not pose any additional unacceptable risks to operators, bystanders and the wider environment, and hence they granted an approval (Health and Safety Executive, 2012).



Safety assessments such as those carried out on acetamiprid are subject to periodic reviews to ensure that conditions of use and any restrictions in approval are in line with the latest internationally peer reviewed evidence and experience. Any changes in the conditions of use specified by the Health and Safety Executive as a result of this process would have to be fully complied with by the forest industry.

Forest Research, and the wider forest industry, relies on this independent, expert assessment of safety, and the summary of the risks involved in using acetamiprid presented in this Report is based largely upon the assessments published by the European Commission (European Commission, 2004a; 2017a), and on the conditions of use specified in the approved product label for Gazelle SG<sup>®</sup> by the Health and Safety Executive (Certis, 2017; Health and Safety Executive, 2017b). These bodies should be contacted directly if further detail of the data, studies and methodology they have used to come to their conclusions is required.

# 6.3. Some hazard and risk characteristics of acetamiprid

Multiple studies have shown that acetamiprid has a lower impact on bee species than other neonicotinoids, and to date it has not been linked to bee decline (e.g. Lundin *et al.*, 2015). It is 400 times less toxic to bees than cypermethrin (based on a contact LD<sub>50</sub> to bees of 0.02 µg per bee for cypermethrin and 8.1 µg per bee for acetamiprid; where 1 µg = 0.000001 g) (European Commission, 2004a; 2005). When acetamiprid is formulated as the product Gazelle SG<sup>®</sup> and used according to the conditions of use on the product label, it is not classified by the Health and Safety Executive as toxic to bees (Certis, 2017).

In common with most insecticides the active ingredient acetamiprid could, in theory, be toxic to bees if it were to be ingested by them. However, once the sprayed product has dried on the planted trees, it is likely that bees would have to ingest parts of the plant to have significant exposure to the insecticide. Since they do not feed on trees and are unlikely to land on the tree stems or foliage when foraging, cross contamination is therefore highly unlikely. No deliberate spraying of surrounding soil or vegetation should take place, so that even if other flowering plants that bees might forage from for nectar and pollen establish on a clearfell site, these plants would not be sprayed, although there is a theoretical risk of misapplication causing spray drift onto these flowering plants. Similarly, no spraying of any small puddles of water that are on the restock site should take place, and there will be little water retained on tree needles, so bees foraging for water are also very unlikely to come in to contact with acetamiprid from treated trees. Honey bees are also unlikely to visit treated trees while sourcing resin to make propolis since they are more likely to obtain resin from the much larger source associated with any nearby mature trees.



The product Gazelle SG<sup>®</sup>, is classified by the Health and Safety Executive as harmful to humans if swallowed (Certis, 2017). Acetamiprid has an acute reference dose (ARfD) of 25 µg per kg of bodyweight (European Commission, 2017a). The acute reference dose is defined as the amount that can be ingested over a short period of time, usually during one meal or one day, without appreciable health risk (European Commission, 2001). This implies that an average person weighing 70 kg could in theory consume 1,750 µg (0.00175 g) of acetamiprid with the likelihood that there would be no short or long term effects on their health. Nevertheless, the aim should be to prevent any exposure to users of the forest, or those living in or close to it. This can be achieved by following normal good working practices such as carefully targeting the spray onto the foliage and stems of small trees, erecting warning signs to warn users to avoid the plants until they are dry, and adopting adequate no-spray buffer zones.

The maximum permitted residue level of acetamiprid on foodstuffs is up to 2 mg (0.002 g) of acetamiprid per kg of fruit (European Commission, 2017b). However under the European Commission Drinking Water Directive there is an even more stringent maximum permissible limit of only 0.1  $\mu$ g l<sup>-1</sup> (i.e. 0.0000001 grams in 1 litre of water) for individual pesticides in tap water in the UK (European Commission, 1998).

The risk to soil living organisms from acetamiprid is low (European Commission 2004a; EFSA 2016). Acetamiprid is not persistent in soil and rapidly degrades by aerobic metabolism to non-toxic metabolites, with a  $DT_{50}$  of 1-5 days (European Commission, 2004a), but it is highly soluble in water and weakly absorbed by the soil, making it potentially susceptible to run-off following application.

The Health and Safety Directorate classify the product Gazelle SG<sup>®</sup> as potentially very toxic to aquatic life with long lasting effects if it is misused, because although it is relatively safe to fish, it can readily kill the early life stages of aquatic insects. However, provided it is used as directed on the approved product label to prevent water contamination, the Health and Safety Executive have concluded that Gazelle SG<sup>®</sup> does not pose an unacceptable risk of harm to the wider environment (Certis, 2017). To help prevent water contamination, one important restriction and condition of their approval for Gazelle SG<sup>®</sup> is the use of 1 metre wide no-spray buffer areas separating treated trees and all watercourses. However, to further minimise the risk of the chemical reaching streams, the UK Forestry Standard extends this level of protection by stipulating the precautionary use of much wider buffers than the 1 metre minimum legal requirement (Forestry Commission, 2017). Recommended buffer widths for all pesticide applications in forestry, which have been adopted in forests being managed to the UK Forestry Standard extends the independently verified in those estates certified via the UK Woodland Assurance Standard (UKWAS, 2018), are as follows:-

 For permanent watercourses with a channel < 2 metres wide – a minimum 10 metre wide unsprayed buffer zone.



- For watercourses > 2 metres wide, and for lakes, reservoirs, large ponds and wetlands a minimum 20 metre wide unsprayed buffer zone.
- Around abstraction points for public and private water supplies, such as springs, boreholes, wells and surface water intakes – a minimum 50 metre wide unsprayed buffer zone.

A permanent watercourse is defined as a stream or river that is delineated on a 1:10,000 Ordnance Survey map, or an open drain that flows directly into a stream. Drains that are separated from watercourses by an adequate buffer area are excluded from this definition. Boggy source areas for streams, even if they are dry at the time of pesticide application, should be incorporated within buffer areas. Before spraying pesticides, checks should be made to ensure that any drainage channels in the area to be treated do not discharge directly into watercourses, and where they do, buffer areas should be extended along these. No-spray buffer zones should also not be used for mixing and filling.

The UK Forestry Standard (Forestry Commission, 2017) includes a number of other water guidelines aimed at protecting the water environment from pesticide use, covering the storage, transport, disposal, handling and application of pesticides. Some of the guidelines have legal status in Scotland as General Binding Rules (SEPA, 2017), while related aspects of good practice such as avoiding spraying in windy conditions to minimise drift, having an emergency plan for dealing with spillages, and safe pesticide disposal, are detailed in the relevant Pesticides Codes of Practice (DEFRA, 2006; Scottish Executive, 2006), which all spray operators are legally obliged to comply with.

If these measures are followed, including the enhanced precautionary buffer zones outlined in the UK Forestry Standard, it is highly unlikely that acetamiprid use in forests will result in any contamination of watercourses, and the risk of exposure of the public to acetamiprid is likely to no greater than that posed by agricultural, horticultural and home garden use, or the risk posed of exposure from residues on food.

Forest Research have recently undertaken a detailed study to specifically check the risk assessment for water by monitoring the downstream effects of planting out acetamiprid pre-treated trees or through using acetamiprid in top-up sprays as described in this Report. The study site was deliberately selected to pose a high risk of chemical run-off, being a high elevation headwater catchment, with poorly drained soils and which was subject to heavy rainfall. The results from this research showed that the usual, good forestry practice measures employed, as described in this Report, were effective in preventing any contamination of stream waters by acetamiprid (Thomas *et al.*, in prep.).



Alice Holt Lodge Farnham Surrey GU10 4LH, UK Tel: 0300 067 5600

info@forestresearch.gov.uk www.forestresearch.gov.uk Northern Research Station Roslin Midlothian EH25 9SY, UK Tel: 0300 067 5900 Forest Research in Wales Environment Centre Wales Deiniol Road Bangor Gwynedd LL57 2UW, UK

Tel: 0300 067 5774

Forest Research will consider all requests to make the content of our documents available in alternative formats. Please send any such requests to: research.info@forestresearch.gov.uk © Crown copyright 2022